



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06621 - Goat Anti-CD274 / PD-L1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CD274 antigen, PD-L1, PDCD1LG1, B7-H, B7H1, PDL1, PDCD1L1, programmed cell death 1 ligand 1, PDL1, HGNC:17635, CD274, CD274 molecule, MGC142294, MGC142296

Official Symbol: CD274

Accession Number(s): NP_054862.1; NP_001254635.1

Human GeneID(s): [29126](#)

Important Comments: This antibody is expected to recognize reported isoforms a and b (NP_054862.1; NP_001254635.1) only.

Immunogen

Peptide with sequence CKKQSDTHLEET, from the C Terminus of the protein sequence according to NP_054862.1; NP_001254635.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 50kDa band observed in Human Heart lysates and in lysates of cell line U2OS (calculated MW of 33.3kDa according to 1NP_054862.1). This band was successfully blocked by incubation with the immunizing peptide and is routinely observed by other sources. Recommended concentration: 0.01-0.1µg/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in th membrane and cytoplasm of U2OS and A431 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of Jurkat cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

Specific References

This antibody (previous batch) has been successfully used in Western blot on Human:

Guozhi Xia, Xiaopu Zheng, Xinye Yao, Xiaowei Yao, Zhongwei Liu, Junkui Wang.


Expression of programmed cell death-1 and its ligand B7 homolog 1 in peripheral blood lymphocytes from patients with peripartum cardiomyopathy.

Clin Cardiol. 2016 Dec 27.

PMID: 28026044

This antibody (previous batch) has been successfully used in Western blot and IHC on Human:


Chen J, Li G, Meng H, Fan Y, Song Y, Wang S, Zhu F, Guo C, Zhang L, Shi Y.
Upregulation of B7-H1 expression is associated with macrophage infiltration in
hepatocellular carcinomas.
Cancer Immunol Immunother. 2012 Jan;61(1):101-8.
PMID: 21853301




EB06621 (0.01 μ g/ml) staining of Human Heart lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.




EB06621 (0.1 μ g/ml) staining of U2OS cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB06621 Immunofluorescence analysis of paraformaldehyde fixed AA431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06621 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06621 Flow cytometric analysis of paraformaldehyde fixed Jurkat cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.