



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06859 - Goat Anti-HMGA1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HMGA1, high mobility group AT-hook 1, HGNC:5010, HMG-R, HMGIY, MGC12816, MGC4242, MGC4854, OTTHUMP00000039618, high-mobility group (nonhistone chromosomal) protein isoforms I and Y, nonhistone chromosomal high-mobility group protein HMG-I/HMG-Y, HMGA1A, OTTHUMP00000016223, OTTHUMP00000016224

Official Symbol: HMGA1

Accession Number(s): NP_665906.1; NP_665908.1; NP_002122.1; NP_665909.1; NP_665912.1; NP_665910.1

Human GeneID(s): [3159](#)

Important Comments: This antibody is expected to recognise isoform a (also called HMG-I; NP_665906.1; NP_665908.1) and isoform b (also called HMG-Y; NP_002122.1; NP_665909.1; NP_665910.1; NP_665912.1).

Immunogen

Peptide with sequence C-LASKQEKDGTEK, from the internal region of the protein sequence according to NP_665906.1; NP_665908.1; NP_002122.1; NP_665909.1; NP_665912.1; NP_665910.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary experiments gave an approx 50kDa band in Human Testis and Ovary lysates after 0.3µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 11.7kDa according to isoform a (also called HMG-I; NP_665906.1; NP_665908.1; NP_665911.1) and 10.7kDa according to isoform b (also called HMG-Y; NP_002122.1; NP_665909.1; NP_665910.1; NP_665912.1). The 50kDa band was successfully blocked by incubation with the immunizing peptide.

IHC: Paraffin embedded Human Placenta and Small Intestine. Recommended concentration: 4µg/ml.

Additional validation: This antibody has been successfully used in qRT-PCR on Human: Cello et al. (2013) PMID: 23545254.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

Specific Reference


This antibody has been successfully used in qRT-PCR on Human:

Di Cello F, Shin J, Harbom K, Brayton C.


Knockdown of HMGA1 inhibits human breast cancer cell growth and metastasis in immunodeficient mice.

Biochem Biophys Res Commun. 2013 Apr 26;434(1):70-4.

PMID: 23545254



EB06859 (4µg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB06859 (4µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.