



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07776 - Goat Anti-AIP / ARA9 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AIP, ARA9, aryl hydrocarbon receptor interacting protein, FKBP16, FKBP37, XAP2, HBV-X associated protein, aryl hydrocarbon receptor-interacting protein

Official Symbol: AIP

Accession Number(s): NP_003968.1

Human GeneID(s): [9049](#)

Non-Human GeneID(s): 11632 (mouse), 282827 (rat)

Immunogen

Peptide with sequence C-EARIRQKDEEDKAR, from the C Terminus of the protein sequence according to NP_003968.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 37kDa band observed in lysates of cell line NIH/3T3 (calculated MW of 37.7kDa according to Human NP_003968.1 and 37.6kDa according to mouse NP_057875.1). In transfected HEK293 transiently expressing ARA9 a band of approx. 42kDa is observed. This band is not observed in the non-transfected HEK293.


Recommended concentration: 0.02-0.06µg/ml.

IHC: Paraffin embedded Human Placenta. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Mouse, Human


Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07776 (0.02 μ g/ml) staining of NIH/3T3 cell lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing ARA9 (RC211157) and probed with EB07776 (mock transfection in first lane), tested by Origene.



EB07776 (5µg/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.