



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10690 - Goat Anti-GNB2L1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** GNB2L1, guanine nucleotide binding protein (G protein), beta polypeptide 2-like 1, Gnb2-rs1, H12.3, HLC-7, PIG21, RACK1, OTTHUMP00000223930, Receptor for Activated C Kinase 1, cell proliferation-inducing gene 21 protein, guanine nucleotide-binding protein subunit beta-2-like 1, guanine nucleotide-binding protein subunit beta-like protein 12.3, human lung cancer oncogene 7 protein, lung cancer oncogene 7, proliferation-inducing gene 21, protein homologous to chicken B complex protein, guanine nucleotide binding, receptor of activated protein kinase C 1

**Official Symbol:** RACK1

**Accession Number(s):** NP\_006089.1

**Human GeneID(s):** [10399](#)

**Non-Human GeneID(s):** 14694 (mouse), 83427 (rat)

### Immunogen

Peptide with sequence NTLGVCKYTVQDESH, from the internal region of the protein sequence according to NP\_006089.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.


**Western blot:** Approx 33kDa band observed in Human Lymph Nodes, in Mouse and Rat Spleen lysates and in lysates of many cell lines (calculated MW of 35.1kDa according to NP\_006089.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Small Intestine. Recommended concentration: 3.75µg/ml.


### Species Reactivity

**Tested:** Human, Mouse, Rat


**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB10690 (0.1 $\mu$ g/ml) staining of Human Lymph Node (A), Mouse Spleen (B) and Rat Spleen (C) lysates (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB10690 (0.1 $\mu$ g/ml) staining of NIH3T3 (A), HeK293 (B), HeLa (C), A431 (D), A549 (E), MCF7 (F), Jurkat (G), K562 (H) lysates (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB10690 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.