



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12354 - Goat Anti-syndecan 1 / CD138 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SDC1, syndecan 1, CD138, SDC, SYND1, syndecan, CD138 antigen, heparan sulfate proteoglycan fibroblast growth factor receptor, syndecan proteoglycan 1, syndecan-1

Official Symbol: SDC1

Accession Number(s): NP_002988.3

Human GeneID(s): [6382](#)

Important Comments: Reported variants represent identical protein: NP_001006947.1, NP_002988.3.

Immunogen

Peptide with sequence C-REQEATPRPRETTQ, from the internal region of the protein sequence according to NP_002988.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 37kDa band observed in preliminary testing of HepG2, cell lysate after 1µg/ml of antibody staining (calculated MW of 32.5kDa according to NP_002988.3). Primary incubation 1 hour at room temperature.

Species Reactivity

Tested:

Expected from sequence similarity: Human