



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12715 - Goat Anti-DC-SIGN / CD209 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CD209, CD209 molecule, CDSIGN, CLEC4L, DC-SIGN, DC-SIGN1, C-type lectin domain family 4 member L, C-type lectin domain family 4, member L, CD209 antigen, HIV gpl20-binding protein, dendritic cell-specific ICAM-3-grabbing non-integrin 1, dendritic cell-specific intracellular adhesion molecules (ICAM)-3 grabbing non-integrin

Official Symbol: CD209

Accession Number(s): NP_066978.1; NP_001138368.1; NP_001138369.1; NP_001138365.1; NP_001138366.1; NP_001138367.1; NP_001138371.1

Human GeneID(s): [30835](#)

Important Comments: This antibody is expected to recognize all reported isoforms.

Immunogen

Peptide with sequence C-NRGEPNNVGEED, from the internal region of the protein sequence according to NP_066978.1; NP_001138368.1; NP_001138369.1; NP_001138365.1; NP_001138366.1; NP_001138367.1; NP_001138371.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 50kDa band observed in Human Bone Marrow lysates (calculated MW of 45.8kDa according to NP_066978.1). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Placenta. Recommended concentration: 6µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human



EB12715 (0.3 μ g/ml) staining of Human Bone Marrow lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB12715 (6 μ g/ml) staining of paraffin embedded Human Placenta. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB12715 Negative Control showing staining of paraffin embedded Human Placenta, with no primary antibody.