

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





BrdU (bromodeoxyuridine). Sheep Polyclonal Antibody

BACKGROUND

Bromodexyuridine (BrdU) is a thymidine analog which is selectively incorporated into the DNA of proliferating cells to provide a marker for the DNA being replicated. The number of proliferating cells can then be detected in cell lysates, tissue sections or suspensions using an antibody specific for the BrdU. Previous methods of detecting DNA included the use of [3H]-thymidine which would be incorporated into the DNA and could then the DNA could be quantified by autoradiography or scintillation counting. These methods are more difficult and require more cleanup due to the radioactive material. An immunohistochemical assay provides a much simpler and cleaner method for detecting DNA in cells. antibody is also useful for detecting proliferating cells by flow cytometry or immunofluorescence staining.

ORDERING INFORMATION

CATALOG NUMBER

A205P

SIZE

 $250 \mu g$

FORM

Unconjugated

HOST/CLONE

Sheep

FORMUL ATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION

See vial for concentration

ISOTYPE

IgG

APPLICATIONS

Western Blot, Immunoprecipitation, Immunohistochemistry

SPECIES REACTIVITY


ACCESSION NUMBER


Not Applicable, Not Applicable

IMMUNOGEN

Bromodeoxyuridine conjugated to *Helix Pomatia* Haemocyanin.

Activity assay using BrdU antibody (Cat. No. A205P) against BrdU labeled BGG (left). Immunoprecipiation assay using BrdU antibody on BrdU, 5-methyl cytosine (5-Me) and control (No Ag).





COMMENTS

Applications: Western blotting: 25 -100 μg/ml; Immunoprecipitation: 25-100 μg/ml; Immunohistochemistry: 25-100 μg/ml The antibody has been tested using immunoprecipitation against 5-methyl cytosine (5-MeC) and bromodeoxyuridine (BrdU) or a control (no antigen). At a concentration of 25 ug/ml, this product demonstrates 8-fold higher reactivity with 5-MeC versus BrdU. For best results, use product at 25-100μg/ml. Note: Denature DNA sample first so that bases are accessible to anti BrdU antibody.

PURIFICATION

Ammonium Sulfate Precipitation

SHIP CONDITIONS

Room Temperature

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

Stability

Products are stable for one year from purchase when stored properly

REFERENCES

- 1. Nakamura, S., et al. Application of bromodeoxyuridine (BrdU) and anti-BrdU monoclonal antibody for the in vivo analysis of proliferative characteristics of human leukemia cells in bone marrows. Oncology 1991, 48, 285-289
- 2. Wilson, G., Cell kinetic studies using a monoclonal antibody to bromodeoxyuridine. Methods Mol. Biol. 1998, 80, 255-266
- 3. Gray, J. (Ed), Special Issue: Monoclonal antibodies against bromodeoxyuridine Cytometry 1985, Vol. 6(6)