



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



A431 cells Control Lysate

BACKGROUND

Total cell lysates are useful as both positive and negative controls in immunoblotting. A431, a human epidermoid carcinoma cell line, and the EGF-stimulated A431 lysates are used as negative and positive controls, respectively, when studying the phosphorylation cascade initiated by ligand binding to receptor tyrosine kinases. Both A431 and normal human fibroblasts are valuable positive controls for antibodies to proteins expressed in these cells. The mouse fibroblast cell line 3T3 and its RSV-transformed counterpart serve as negative and positive controls, respectively, for antibodies to proteins that are phosphorylated as a result of transformation by RSV (Rous sarcoma virus). Madin-Darby bovine kidney (MDBK) cell lysate displays proteins expressed in bovine kidney and PC12 (rat pheochromocytoma) cell lysate can be expected to mirror the protein expression of cells and tissues of neuroectodermal origin. Likewise, Jurkat cells express the proteins unique to human T-cells. Rat brain is a widely used positive control for those proteins that are exclusively expressed in brain. At right is an example of the use of positive and negative control lysates in western blotting. The blot of 3T3 cell lysate (lane 1) and RSV-transformed 3T3 (lane 2) was probed with anti-phosphotyrosine peroxidase conjugate to reveal those proteins that are tyrosine-phosphorylated in response to transformation by

ORDERING INFORMATION

CATALOG NUMBER

X1002

SIZE

100 μ l

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in Laemmli electrophoresis sample buffer. Samples need to be boiled for 3-5 minutes, before loading the gel.

SHIP CONDITIONS

Room Temperature

STABILITY

Reagents are stable for the period shown on the vial label when stored properly

COMMENTS

A volume of 10-25ul is suggested for immunoblotting.