



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Collagen Substrate, FITC Conjugate. Purified Protein
Extracted from bovine skin

BACKGROUND

FITC-labeled bovine type I is an excellent substrate for examining collagenase activity. This labelled substrate is highly purified and telopeptide free and is supplied lyophilized. To minimize background levels in collagenase assays, FITC-labeled collagen has been enzymatically pre-treated and further purified by ion-exchange chromatography.

FITC-labeled collagen can also be used as a substrate for cell culture. Collagen degradation products in the culture supernatants can be directly determined by measured the fluorescence at 520 nm (Emission)/490 nm (Excitation).

ACTIVITY

Used for the assay of mammalian collagenase activity. To heat denature collagen for preparing a substrate for gelatinase, transfer collagen solution to a brown glass bottle and heat to 80°C.

PURITY

>95%

REFERENCES

1. Terato, K., et al. "A rapid assay method of collagenase activity using 14C-labeled soluble collagen as substrate." *Biochim. Biophys. Acta* 1976, 445, 753-762
2. Sellers, A., et al. "Evidence that latent collagenases are enzyme-inhibitor complexes." *Biochem. J.* 1977, 163, 303-307
3. Shinkai, H., et al. "A complex of collagenase with low molecular weight inhibitors in the culture medium of embryonic chick skin explants." *J. Biochem. (Tokyo)* 1977, 81, 261-263
4. Shinkai, H. & Nagai, Y., "A latent collagenase from embryonic human skin explants." *J. Biochem. (Tokyo)* 1977, 81, 1261-1263
5. Murawaki, Y., et al. "Serum collagenase activity in patients with chronic liver disease." *J. Hepatol.* 1993, 328-334

ORDERING INFORMATION

CATALOG NUMBER

X1098

SIZE

10 mg

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as a 1 mg/ml solution in 0.01M acetic acid.

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STABILITY

Stable in acidic buffer when stored at -20°C, however slowly degrades when

CONCENTRATION

See vial for concentration

SOURCE

Bovine skin