



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Sphingosine 1 Phosphate 4 Receptor (NT) (EDG6) NT. Rabbit Polyclonal Antibody

BACKGROUND

EDG6 (S1P₄) belongs to a family of G-protein coupled receptors whose ligands are lysophospholipids. The ligand for EDG6 (S1P₄) is sphingosine 1-phosphate. There are 8 known members of the EDG receptor family and they are implicated in mediating growth related effects such as induction of cellular proliferation, alterations in differentiation and survival and suppression of apoptosis. They also evoke cellular effector functions that are dependent on cytoskeletal responses such as contraction, secretion, adhesion and chemotaxis. EDG receptors are developmentally regulated and differ in tissue distribution. They couple to multiple types of G proteins to signal through ras and MAP kinase, rho, phospholipase C and several protein tyrosine kinases. EDG6 (S1P₄) is expressed in lymphoid and hematopoietic tissue and in the lungs.

IMMUNOGEN

Synthetic peptide derived from the N terminal of the EDG6 (S1P₄) receptor

POSITIVE CONTROL/TISSUE EXPRESSION

Cos cells transfected with EDG6 (S1P₄) protein

COMMENTS

Detects EDG6 (S1P₄) receptors at concentration of 5-10 µg/ml by Western blot using a human EDG6 (S1P₄) receptor transfected cell line. Detects an approximately 45 kDa band in Cos cells transfected with full length human EDG6 (S1P₄). Due to low expression of EDG receptors, we recommend use of Pierce Femto Signal substrate for western blot development.

ORDERING INFORMATION

CATALOG NUMBER
X1593P

SIZE
100 µg
FORM
Unconjugated

HOST/CLONE
Rabbit

FORMULATION
Provided as solution in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION
See vial for concentration

ISOTYPE
IgG

APPLICATIONS
Western Blot

SPECIES REACTIVITY
Human

ACCESSION NUMBER
O95977, Human

PURIFICATION

Ammonium Sulfate Precipitation

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

1. Yamazaki, Y., et al. "EDG-6 as a putative sphingosine 1-phosphate receptor coupling to Ca(2+) signaling pathway." *Biochem. Biophys. Res. Commun.* 2000, 268, 583-589.
2. Van Brocklyn, J.R., et al. "Sphingosine 1-phosphate is a ligand for the G protein-coupled receptor EDG-6." *Blood* 2000, 95, 2624-2629.
3. Takuwa, Y., et al. "Subtype-specific, differential activities of the EDG family receptors for sphingosine 1-phosphate, a novel lysophospholipid mediator." *Mol. Cell Endocrinol.* 2001, 177, 3-11.