



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



SPARC Blocking Peptide

BACKGROUND

SPARC is a key factor in cell-matrix interactions and possibly tumour aggressiveness. The SPARC gene, which encodes a multifunctional glycoprotein with roles in tissue development, remodelling and fibrosis. A regulator of cell-extracellular matrix (ECM) interactions, SPARC represents a major factor in the ECM remodelling occurring during tumour invasion. *in silico* analysis reveals 4 UTR-SNPs located in the 3'-UTR of the SPARC gene, corresponding to 1474 g a, 1551 g c, 1922 t g and 2072 c t changes, which are significantly associated with tumoral state of the tissue. Of all hits, the 2072 SPARC polymorphism had the best association with cancer.

SPARC therefore is a gene involved in a number of diseases including rheumatoid arthritis, scleroderma, tumor development and metastasis. SPARC variants have been detected in tumour samples of patients with acute myeloblastic leukemia (AML).

ORDERING INFORMATION

CATALOG NUMBER

X1855B

SIZE

50 μ g

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STABILITY

Products are stable for one year from purchase when stored properly

COMMENTS

Blocking peptide for use with SPARC antibodies (Cat. Nos. X1860P & X1867P).

INSTRUCTIONS

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 μ l or 10 μ g of Exalpha's rabbit anti-SPARC (Cat. No. X1860P or X1867P) is added to 10 μ g of blocking peptide for a total volume of 20 μ l. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).