



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FHL2. Rabbit Antigen Immunoaffinity Purified Polyclonal
Four-and-a-half LIM only protein 2, FHL2

BACKGROUND


Sphingosine kinase-1 (SK1) is a key enzyme that catalyzes the phosphorylation of sphingosine to sphingosine-1-phosphate (S1P). SK1, and its product S1P, regulate numerous and diverse biological functions, including cell growth, differentiation, proliferation, and apoptosis. S1P also plays a central role in cardiac development and ischemic preconditioning. FHL2 (SLIM3) has been identified as a SK1-interacting protein in mammalian cardiomyocytes. FHL2, but not FHL1 or FHL3, interacts with SK1, and FHL2 colocalizes with SK1 in the cytoplasm of these cells. The interaction of FHL2 with SK-1 involves the C-terminal portion of SK1. Overexpression of FHL2 has been found to attenuate the activity and antiapoptotic effects of SK1. Endothelin-1, a potent survival factor in cardiomyocytes, inhibits FHL2-SK1 association and increases SK1 activity. FHL2 is therefore a novel inhibitor of SK1 activity in cardiomyocytes and targeting FHL2 may prevent myocardial apoptosis through activation of SK1.

A transcriptional coactivator, FHL2 is also known to form complexes consisting of Proline-, glutamic acid-, and leucine-rich protein-1 (PELP1), FHL2, and androgen receptors (AR) in prostate cancer cells, perhaps functioning as a molecular adaptor with PELP1 in the progression of prostate cancer.

IMMUNOGEN

Synthetic peptide derived from the human FHL2 protein

Western blot using Exalalpha's X1857P, rabbit polyclonal at 0.5 ug/ml on HeLa cell extract (10 ug/lane). Blots were developed with goat anti-rabbit Ig (1:75k) and Pierce's Supersignal West Femto system.



ORDERING INFORMATION

CATALOG NUMBER
X1857P

SIZE
50 µg
FORM
Affinity Purified

HOST/CLONE
Rabbit

FORMULATION
Provided as ligand affinity purified antibody in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION
Lot specific, see vial for details

ISOTYPE
IgG

APPLICATIONS
Western Blot

SPECIES REACTIVITY
Human

ACCESSION NUMBER
Human CAG33718

POSITIVE CONTROL/TISSUE EXPRESSION

Antibody has been tested on HeLa cell lysates.

COMMENTS

Antibody can be used for Western blotting (0.5-2 μ g/ml). Also binds ABB73038, FHL2 isoform 5 with MW of 389 a.a. (~42kDa). Optimal concentration should be evaluated by serial dilutions.

PURIFICATION

Antigen Immunoaffinity Purification

SHIP CONDITIONS

Ship on dry ice, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -70°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

- 1: Nair SS, et al. PELP1/MNAR enhances androgen receptor functions through LIM-only coactivator FHL2. Mol Endocrinol. 2006 Dec 27
- 2: Kudo LC, et al. Genetic Analysis of Anterior-Posterior Expression Gradients in the Developing Mammalian Forebrain. Cereb Cortex. 2006 Dec 5;
- 3: Kahl P, et al. Androgen receptor coactivators lysine-specific histone demethylase 1 and four and a half LIM domain protein 2 predict risk of prostate cancer recurrence. Cancer Res. 2006 Dec 1;66(23):11341-7.
- 4: Campo-Fernandez B, et al. Identification of the FHL2 Transcriptional Coactivator as a New Functional Target of the E7 Oncoprotein of Human Papillomavirus Type 16. J Virol. 2007 Jan;81(2):1027-32. Epub 2006 Nov 8.
- 5: Govoni KE, et al. Disruption of four-and-a-half LIM 2 decreases bone mineral content and bone mineral density in femur and tibia bones of female mice. Calcif Tissue Int. 2006 Aug;79(2):112-7. Epub 2006 Aug 15.
- 6: Sun J, et al. FHL2/SLIM3 decreases cardiomyocyte survival by inhibitory interaction with sphingosine kinase-1. Circ Res. 2006 Sep 1;99(5):468-76. Epub 2006 Aug 3.

PRODUCT SPECIFIC REFERENCES