



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Sphingosine Kinase 1, Mouse Blocking Peptide

BACKGROUND

Sphingosine kinase 1 catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (spp)(sphingosine + ATP = sphingosine 1-phosphate + ADP), a lipid mediator with both intra- and extracellular functions. also acts on d-erythro-sphingosine and to a lesser extent sphinganine, but not other lipids, such as d,l-threo-dihydrosphingosine, n, n-dimethylsphingosine, diacylglycerol, ceramide, or phosphatidylinositol. The long form of sphingosine kinase 1, an N-terminal extension, is responsible for FceRI-mediated calcium mobilization and may play a role in other ligand mediated signaling events.

ORDERING INFORMATION

CATALOG NUMBER

X1878B

SIZE

50 μ g

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

SHIP CONDITIONS

Ship on dry ice, freeze upon arrival

STABILITY

Products are stable for one year from purchase when stored properly

COMMENTS

Blocking peptide for use with mouse sphingosine kinase 1 antibody (Cat. No. X1882P).

INSTRUCTIONS

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 μ l or 10 μ g of Exalalpha's rabbit anti-mouse Sphingosine Kinase 1 (Cat. No. X1882P) is added to 10 μ g of blocking peptide for a total volume of 20 μ l. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).