



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Sphingosine Kinase 2, Long Form, Human. Rabbit Antigen Immunoaffinity Purified Polyclonal
Sphk2-L, N-terminal-extended Sphingosine Kinase 2

BACKGROUND

Rat sphingosine kinase 1 catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (spp)(sphingosine + ATP = sphingosine 1-phosphate + ADP), a lipid mediator with both intra- and extracellular functions. also acts on d-erythro-sphingosine and to a lesser extent sphinganine, but not other lipids, such as d,l-threo-dihydrosphingosine, n, n-dimethylsphingosine, diacylglycerol, ceramide, or phosphatidylinositol. The long form of sphingosine kinase 1, an N-terminal extension, is responsible for FcεRI-mediated calcium mobilization and may play a role in other ligand mediated signaling events.

ORDERING INFORMATION

CATALOG NUMBER
X1880P

SIZE
10 Miniblots

FORM
Affinity Purified

HOST/CLONE
Rabbit

FORMULATION
Provided as ligand affinity purified antibody in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION
Lot specific, see vial

ISOTYPE
IgG

APPLICATIONS
Western Blot, ELISA


SPECIES REACTIVITY
Human

ACCESSION NUMBER
Human Q9NRA0

IMMUNOGEN

Synthetic peptide derived from the N-terminal extension of the Sphingosine Kinase 2-L protein.

Western blot of endogenous sphingosine kinase 2 LF detection in Jurkat cell lysate (4 ug/lane) using X1880P (1 ug/ml) developed using anti-rabbit HRP (1:75K) and Pierce Super Signal west Femto.



POSITIVE CONTROL/TISSUE EXPRESSION

Jurkat lysate used as positive control.

COMMENTS

Antibody can be used for Western blotting (1-5 μ g/ml). Optimal concentration should be evaluated by serial dilutions.

PURIFICATION

Antigen Immunoaffinity Purification

SHIP CONDITIONS

Ship on dry ice, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -70°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

1. Okada, T., et al. 'Involvement of N-terminal-extended form of sphingosine kinase 2 in serum-dependent regulation of cell proliferation and apoptosis.' J. Biol. Chem. 2005, 280, 36318-36325

PRODUCT SPECIFIC REFERENCES