



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Adiponectin Receptor 2 Blocking Peptide

BACKGROUND

ADR2 functions as a receptor for globular and full-length adiponectin (APM1), which is an essential hormone secreted by adipocytes that acts as an antidiabetic. ADR2 is also probably involved in metabolic pathways that regulate lipid metabolism such as fatty acid oxidation. ADR2 mediates an increase in AMPK, PPARA ligand activity, fatty acid oxidation and glucose uptake by adiponectin and has some intermediate-affinity receptor activity for both globular and full-length adiponectin. ADR2 may form homo and heteromultimers.

ORDERING INFORMATION

CATALOG NUMBER

X2051B

SIZE

50 μ g

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STABILITY

Products are stable for one year from purchase when stored properly

COMMENTS

For use with Adiponectin Receptor 2 polyclonal antibody (Cat. No. X1699P).

INSTRUCTIONS

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 μ l or 10 μ g of Exalpa's rabbit anti-Adiponectin Receptor 2 (Cat. No. X1699P) is added to 10 μ g of blocking peptide for a total volume of 20 μ l. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).