



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## SMase Activation Associated Factor Blocking Peptide

### BACKGROUND

Novel adaptor protein which mediates the activation of sphingomyelinase by receptors such as tumor necrosis factor receptor type 1 and CD40 and subsequent apoptosis in cardiomyocytes. Inhibition of this factor may provide a novel intervention point for reducing ischemia/reperfusion injury.

### ORDERING INFORMATION

**CATALOG NUMBER**

X2289B

**SIZE**

50  $\mu$ g

**CUSTOMER STORAGE**

Product should be stored at -20°C.  
Aliquot to avoid freeze/thaw cycles

**FORMULATION**

Provided as solution in phosphate buffered saline

**SHIP CONDITIONS**

Ship at ambient temperature, freeze upon arrival

**STABILITY**

Products are stable for one year from purchase when stored properly

### COMMENTS

For use with FXBW7 polyclonal antibodies (Cat. No. X2314P & X2336P).

### INSTRUCTIONS

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10  $\mu$ l or 10  $\mu$ g of Exalpha's rabbit anti- (Cat. No. ) is added to 10  $\mu$ g of blocking peptide for a total volume of 20  $\mu$ l. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).