



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Lass1 (Cross reacts with GDF1) Blocking Peptide

BACKGROUND

LAG1 is a longevity gene, the first such gene to be identified. Originally cloned from the yeast *Saccharomyces cerevisiae*. A close homolog of this gene, LAC1, has been found in the yeast genome. The human homolog of LAG1 has functions in human aging. LAG1 may be involved in neurodegenerative diseases and human aging. Lass1 may be either a bona fide (dihydro)ceramide synthase or a modulator of its activity. When overexpressed in cells is involved in the production of sphingolipids containing mainly one fatty acid donor (N-linked stearyl- (C18) ceramide) in a fumonisin B1-independent manner. Located in the endoplasmic reticulum membrane; LAss1 is a multi-pass membrane protein.

ORDERING INFORMATION

CATALOG NUMBER

X2294B

SIZE

50 μ g

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STABILITY

Products are stable for one year from purchase when stored properly

COMMENTS

For use with Lass1 polyclonal antibodies (Cat. No. X2084P & X2310P).

INSTRUCTIONS

Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37 deg C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 μ l or 10 μ g of Exalpa's rabbit anti-Lass1 (Cat. No. X2084P) is added to 10 μ g of blocking peptide for a total volume of 20 μ l. The mixture is allowed to incubate for 20 minutes at 37 deg C prior to dilution in suitable buffer (for Western blot, etc.).