



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ADAM DEC1. Rabbit Polyclonal Antibody

A disintegrin and metalloproteinase domain-like protein decysin 1, ADAM-like protein decysin 1

BACKGROUND

ADAMDEC1 is expressed in the immune system, by dendritic cells and macrophages. It may play an important role in the control of the immune response and during pregnancy.¹ Due to the partial lack of a disintegrin domain and the total lack of a cysteine-rich domain, this protein has been placed in a novel subclass of the ADAM gene family².

ORDERING INFORMATION

CATALOG NUMBER
X2416P

SIZE

100 µg

FORM

Unconjugated

HOST/CLONE

Rabbit

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION

See vial for concentration

ISOTYPE

IgG

APPLICATIONS

Western Blot, ELISA

SPECIES REACTIVITY

Human


ACCESSION NUMBER

Human O15204

IMMUNOGEN

Synthetic peptide derived from the human ADAMDEC1 protein.

Staining on paraffin embedded normal human tonsil sections. Primary Ab (IgG cut or immunoaffinity purified) at 10 µg/ml. Antigen retrieval used: 10 mM Na Citrate pH 6.0, 10 minutes pressure cooker method., developed with anti rabbit HRP and DAP substrate. Counterstained with methyl blue.



POSITIVE CONTROL/TISSUE EXPRESSION

Expressed highly in the small intestine and appendix, moderately in lymph node, mucosal lining of the colon, thymus, spleen and very weakly in the bone marrow. Predominantly expressed in dendritic cells (DC) of the germinal center. Weakly expressed in monocyte and highly expressed in macrophage. Absent in immature DC.

COMMENTS

Antibody can be used for Western blotting (1-5 μ g/ml). Optimal concentration should be evaluated by serial dilutions.

PURIFICATION

Ammonium Sulfate Precipitation

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Reagents are stable for the period shown on the vial label when stored properly

REFERENCES

1. UniProtKB/Swiss-Prot Acession # O15204
2. Bates, E.E. et. atl., The ADAMDEC1 (decysin) gene structure: evolution by duplication in a metalloprotease gene cluster on chromosome 8p12. Immunogenetics. 2002 May;54(2):96-105. Epub 2002 Mar 16,

PRODUCT SPECIFIC REFERENCES