

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



calcium channel subunit alpha-11. Rabbit Antigen Immunoaffinity Purified Polyclonal

BACKGROUND

Voltage-sensitive calcium channels (VSCC) mediate the entry of calcium ions into excitable cells and are also involved in a variety of calciumdependent processes, including muscle contraction, hormone or neurotransmitter release, gene expression, cell motility, cell division and cell death. T-type channels serve pacemaking functions in both central neurons and cardiac nodal cells and support calcium signaling in secretory cells and vascular smooth muscle. They may also be involved in the modulation of firing patterns of neurons which is important for information processing as well as in cell growth processes. Using co-transfection techniques we investigated the functional effects of each of the gamma subunits on the biophysics of the T-type VSCC encoded by the alpha(11) subunit. This revealed a substantially slowed rate of deactivation in the presence of gamma(2). In contrast, there was no significant corresponding effect of either gamma(3) or gamma(4) on alpha(11) subunit-mediated currents.(2) **ORDERING INFORMATION** CATALOG NUMBER X2716P SIZE 10 Miniblots FORM Affinity Purified HOST/CLONE Rabbit FORMULATION Provided as solution in phosphate buffered saline with 0.08% sodium azide CONCENTRATION See vial for concentration ISOTYPE lgG **APPLICATIONS** Western Blot SPECIES REACTIVITY Human

ACCESSION NUMBER

IMMUNOGEN

Synthetic peptide derived from the rat alpha1I calcium channel conjugated to KLH

POSITIVE CONTROL/TISSUE EXPRESSION

Rat brain lysate

COMMENTS

Antibody can be used for Western blotting (1:400 dilution). Optimal concentration should be evaluated by serial dilutions.

For research use only. Not for use in human diagnostics or therapeutics.

Exalpha Biologicals, Inc. 2 Shaker Road, Bldg. B101 Shirley, MA 01464 Tel: 800.395.1137 Fax: 866.924.5100 www.exalpha.com info@exalpha.com Page 1 of 2 Cat. No. X2716P

PURIFICATION

Antigen Immunoaffiinity Purification

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

1. McRory, J.E. et al "Molecular and functional characterization of a family of rat brain T-type calcium channels." J. Biol. Chem. 276 (6), 3999-4011 (2001).

2. Green, P.J., et al.Kinetic modification of the alpha(1I) subunit-mediated T-type Ca(2+) channel by a human neuronal Ca(2+) channel gamma subunit." J. Physiol. 533(Pt 2):467-78.(2001)

For research use only. Not for use in human diagnostics or therapeutics.