



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

beta-Actin, (cytoplasmic 1) (NT). Rabbit Antigen Immunoaffinity Purified Polyclonal

BACKGROUND

Actins are highly conserved proteins that are involved in cell motility, structure and integrity, processes that are crucial for tissue development and the development of organism. The actin cytoskeleton is one of the principal drivers of cell motility and is capable of responding to complex signaling cascades. Recent evidence suggests that it may play key roles in regulating apoptosis and aging. Beta actin is one of six different actin isoforms which have been identified. Like GAPDH, b-actin is constitutively expressed at high levels in almost all tissues and cell lines making it ideal for use as a loading control marker in immunoblots.

ORDERING INFORMATION

CATALOG NUMBER
X2742P

SIZE
100 μ g

FORM
Affinity Purified

HOST/CLONE
Rabbit

FORMULATION
Provided as solution in phosphate buffered saline with 0.08% sodium azide

CONCENTRATION
See vial for concentration

ISOTYPE
Ig

APPLICATIONS
Western blot, Immunohistochemistry


SPECIES REACTIVITY
Human

ACCESSION NUMBER
Human P60709

IMMUNOGEN

Synthetic peptide derived from the N-terminal of the beta-actin protein.

Immunohistochemistry image using beta-Actin (NT) antibody (Cat. No. X2742P) on normal human liver tissue sections at 10 μ g/ml.



POSITIVE CONTROL/TISSUE EXPRESSION

Human liver tissue

COMMENTS

Antibody can be used for Western blotting (1-5 µg/ml) and immunohistochemistry on paraffin-embedded tissues (10 µg/ml), Optimal concentration should be evaluated by serial dilutions.

PURIFICATION

Antigen Immunoaffinity Purification

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STORAGE CUSTOMER

Product should be stored at -20°C. Aliquot to avoid freeze/thaw cycles

STABILITY

Products are stable for one year from purchase when stored properly

REFERENCES

1. Ponte P., Ng S.Y., Engel J., Gunning P., Kedes L., Evolutionary conservation in the untranslated regions of actin mRNAs: DNA sequence of a human beta-actin cDNA. Nucleic Acids Res. 12:1687-1696(1984)
2. Nakajima-Iijima S., Hamada H., Reddy P., Kakunaga T., Molecular structure of the human cytoplasmic beta-actin gene: interspecies homology of sequences in the introns. Proc. Natl. Acad. Sci. U.S.A. 82:6133-6137(1985)
3. Procaccio V., Salazar G., Ono S., Styers M.L., Gearing M., Davila A., Jimenez R., Juncos J., Gutekunst C.-A., Meroni G., Fontanella B., Sontag E., Sontag J.-M., Faundez V., Wainer B.H. A mutation of beta -actin that alters depolymerization dynamics is associated with autosomal dominant developmental malformations, deafness, and dystonia. Am. J. Hum. Genet. 78:947-960(2006) [PubMed: 16685646] [Abstract]

PRODUCT SPECIFIC REFERENCES