



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Histone H2a.x Blocking Peptide

BACKGROUND

This peptide is the immunogen used to make Cat # X2753P.

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

ORDERING INFORMATION

CATALOG NUMBER

X2768B

SIZE

50 μ g

CUSTOMER STORAGE

Product should be stored at -20°C.
Aliquot to avoid freeze/thaw cycles

FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

STABILITY

Products are stable for one year from purchase when stored properly

COMMENTS

Optimal concentration should be evaluated by serial dilutions.

INSTRUCTIONS

Use: Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37°C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 μ l or 10 μ g of Exalpha's rabbit anti-Histone H2a.x (Cat. No. X2753P) is added to 10 μ g of blocking peptide for a total volume of 20 μ l. The mixture is allowed to incubate for 20 minutes at 37°C prior to dilution in suitable buffer (for Western blot, etc.).