



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Donkey Chicken IgY, conjugated with Agarose

[exalpha.com/products/donkey-chicken-igy-conjugated-with-agarose-2/DAIgY-AGA-2](https://www.exalpha.com/products/donkey-chicken-igy-conjugated-with-agarose-2/DAIgY-AGA-2)

Catalog number: **DAIgY-AGA-2**

Product Type	Secondary antibodies
--------------	----------------------

Units	2.0 ml bed volume
-------	-------------------

Host	Donkey
------	--------

Source

A donkey was immunized with purified chicken IgY. Antibodies were isolated from immune sera using affinity chromatography and conjugated to N- hydroxysuccinimide (NHS)-activated agarose beads. Beads were washed extensively after the blocking of the residual NHS sites.

Product

Gel; 2 ml settled gel

Product Form: Immunoprecipitation Gel

Formulation: Phosphate buffered saline, pH 7.2 with 0.075% sodium azide

Purification Method: Egg-yolk derived IgY Affinity Purified

Concentration: N/A

Applications

The specific activity of the anti-IgY-agarose was determined by applying 2.7 mg IgY in 24 ml PBS (0.1 mg/ml) to 1.0 ml bed volume of agarose. Two ml aliquots were collected. The total amount of IgY that bound to the column was calculated. For this lot, 1.3 mg IgY bound per ml agarose.

Storage

Product should be stored at 4-8°C. DO NOT FREEZE

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional

information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpa Biologicals accepts no liability for any inaccuracies or omissions in this information.

References

- Hall, Sarah E., Song Luo, Anne E. Hall, and Daphne Preuss, 2005, Differential Rates of Local and Global Homogenization in Centromere Satellites From Arabidopsis Relatives. *Genetics*: 170(4): 1913–1927.
- Song, M-G and M. Kiledjian, 2007. 3' Terminal oligo U-tract-mediated stimulation of decapping. *RNA*, 13:2356-2365.
- Taylor, A.I., H.J. Gould, B.J. Sutton and R.A. Calvert. 2008. Avian IgY Binds to a Monocyte Receptor with IgG-like Kinetics Despite an IgE-like Structure. *J. Biol. Chem.* 283: 16384-16390.
- S-C. Lo and M. Hannink. 2008. PGAM5 tethers a ternary complex containing Keap1 and Nrf2 to mitochondria. *Experimental Cell Research* 314: 1789-1803.
- C. Butkinaree, W.D. Cheung, S. Park, K. Park, M. Barber and G.W. Hart, 2008. Characterization of Beta-N-Acetylglucosaminidase Cleavage by Caspase-3 during Apoptosis. *J. Biol. Chem.*, 283:23557-23566.
- Slawson, C. Lakshmanan, T., Knapp, S. and Hart, G.W., 2008, A Mitotic GlcNAcylation/Phosphorylation Signaling Complex Alters the Posttranslational State of the Cytoskeletal Protein Vimentin. *Mol. Biol. Cell* 19:4130-4140.
- McMichael, Brooke K., Cheney, Richard E., and Beth S. Lee. 2010. Myosin X Regulates Sealing Zone Patterning in Osteoclasts through Linkage of Podosomes and Microtubules. *J Biol. Chem.* 285:9506-9515.
- Hale, C.R., Sonali, M., Elmore, J., Pfister, N., Compton, M., Olson S., Resch, A.M., Glover III, C.V.C., Graveley B.R., Terns, R.M., and M.P. Terns. 2012. Essential Features and Rational Design of CRISPR RNAs that Function with the Cas RAMP Module Complex to Cleave RNAs. *Molecular Cell* 45:292-302.
- Lehman, HL, van Laere SJ, van Golen, CM, Vermeulen PB, Dirix LY and KL van Golen. 2012. Regulation of inflammatory breast cancer cell invasion through Akt1/PKB? phosphorylation of RhoC GTPase. *Mol Cancer Res.* 10:1306-18.
- Cheadle L. and T. Biederer. 2012. The novel synaptogenic protein Farp1 links postsynaptic cytoskeletal dynamics and transsynaptic organization. *J Cell Biol.* 199:985-1001.
- Dai, L, Taylor, MS, O'Donnell, KA and JD Boeke. 2012. Poly (A) Binding Protein C1 is Essential for Efficient L1 Retrotransposition and Affects L1 RNP Formation. *Mol Cell Biol.* 32:4323-4336.
- Joglekar, M, Elbazanti, WO, Weizman, MD, Lehman, HL and van Golen, KL. 2015. Caveolin-1 Mediates Inflammatory Breast Cancer Cell Invasion via the Akt1 Pathway and RhoC GTPase. *J. Cell. Biochem* 116:923-933.
- Zhang, Z., Costa, FC, Tan, EP, Bushue, N, DiTacchio, N, Costello, CE, McComb, ME, Whelan, SA, Petersen, KR, and Slawson, C. 2016. O-Linked N-Acetylglucosamine (O-GlcNAc) Transferase and O-GlcNAcase Interact with Mi2? Protein at the A?-Globin Promoter. *J. Biol Chem* 291:15628-40.
- Balachandran RS, Heighington CS, Starostina NG, Anderson JW, Owen DL, Vasudevan S, Kipreos ET. 2016. The ubiquitin ligase CRL2ZYG11 targets cyclin B1 for degradation in a conserved pathway that facilitates mitotic slippage. *J Cell Biol.* 215(2):151-166.

Safety Datasheet(s) for this product:

EA_Sodium Azide

