



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Fc epsilon R1 alpha antibody [CRA1] (FITC)

Cat No. GTX00852

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Application	WB, ICC/IF, IHC-P, IHC-Fr, FACS, Functional Assay
Reactivity	Human, House musk shrew

Package
50 µg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
WB	Assay dependent
ICC/IF	2-10 µg/ml
IHC-P	2-10 µg/ml
IHC-Fr	2-10 µg/ml
FACS	2-10 µg /ml
Functional Assay	Assay dependent

Note : Titration of IgE-bound fraction of the Fc epsilon R1 alpha using clone CRA1 and CRA2 antibodies.

Not tested in other applications.

Calculated MW 30 kDa. ([Note](#))

Specificity/Sensitivity

The CRA1 (AER37) monoclonal antibody reacts with the Fc epsilon R1 alpha subunit on a region that does not overlap the region of the IgE binding site, thus it does not compete with IgE for the receptor binding. Since the CRA2 (AER24) monoclonal antibody reacts with the IgE binding site on Fc epsilon R1 alpha, it competes with IgE for the receptor binding. Combining the two antibodies, one can quantitatively measure the amounts of the IgE-bound Fc epsilon R1 alpha.

PROPERTIES

Form	Liquid
Buffer	Filter-sterilized PBS (pH 7.4), 50% glycerol
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant extracellular protein of human Fc epsilon R1 alpha; (corresponding to amino acids Met-26-197, where signal peptide is 1-25) Epitope: 26-110 amino acids



For full product information, images and publications, please visit our [website](#).

Date 2020 / 07 / 27 Page 1 of 2

Purification	Purified IgG From culture supernant
Conjugation	Fluorescein isothiocyanate (FITC)
Note	For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).