



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## CD40 antibody [FGK45]

**Cat No. GTX01483**

<b>Host</b>	Rat
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2a
<b>Application</b>	FACS, Activation
<b>Reactivity</b>	Mouse

Reference ( 4 )  
 Package  
 100 µg

## PRODUCT

## Summary

The FGK45 antibody reacts with mouse CD40, a 40-50 kDa glycoprotein member of the tumor necrosis factor receptor (TNFR) family. CD40 is constitutively expressed by antigen presenting cells, including dendritic cells, B cells, and macrophages. The interaction of CD40 and its ligand CD154, expressed on activated T cells, plays an essential role in mediating a broad range of immune and inflammatory responses including T cell dependent immunoglobulin class switching, germinal center formation, and the development of memory B cells. The FGK45 clone is reported to be functionally useful for both in vitro and in vivo activation of CD40 expressing antigen presenting cells. The FGK45 antibody reacts with mouse CD40, a 40-50 kDa glycoprotein member of the tumor necrosis factor receptor (TNFR) family.

## APPLICATION

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
FACS	Assay dependent
Activation	Assay dependent
Inhibition assay	Assay dependent
Not tested in other applications.	

**Calculated MW** 32 kDa. ( [Note](#) )

## PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	10 mM NaH <sub>2</sub> PO <sub>4</sub> (pH 7.2), 150 mM NaCl, 0.09% sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
<b>Concentration</b>	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Purification</b>	Purified by affinity chromatography From tissue culture supernatant
<b>Conjugation</b>	Unconjugated



For full product information, images and publications, please visit our [website](#).

**Note**

For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).