



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# WGA Lectin (Biotin)

**Cat No. GTX01501**
**Application** WB, ELISA, IHC

**Package**

1 mg

**PRODUCT**
**Product Description**

Wheat germ agglutinin (WGA) (MW 36 kDa) is a dimeric, carbohydrate-free protein composed of two identical subunits. The amino acid sequence provides a figure of 21,600, based on 171 amino acids per polypeptide chain. The protein dissociates into monomers in acidic media, pK of the reaction 4. The receptor sugar for WGA is N-acetyl glucosamine (GlcNAc), GlcNAc $\beta$ 14GlcNAc. These structures are present in many serum and membrane glycoproteins, bacterial cell wall peptidoglycans, chitin, cartilage glycosaminoglycans and glycolipids. Native WGA also react with glycoproteins with sialic acid residue. This lectin is useful for the purification of insulin receptors, serum proteins and neuronal tracing. The carbohydrate-binding specificity of WGA has been studied by variety of techniques, such as hapten inhibition of hemagglutination and specific precipitation of glycoconjugate, changes in fluorescence of the lectin (intrinsic) or of chromatogenic ligands (extrinsic), equilibrium dialysis, NMR and x-ray diffraction.

**APPLICATION**
**Specificity/Sensitivity** N-acetyl glucosamine (GlcNAc)

**PROPERTIES**
**Form** Liquid

**Buffer** 10 mM phosphate, 150 mM NaCl, 0.1 mM Calcium chloride (pH 7.6) and 0.05% sodium azide.

**Storage** Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4° C.

**Concentration** 5 mg/ml (Please refer to the vial label for the specific concentration.)

**Conjugation** Biotin

**Note** For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

 For full product information, images and publications, please visit our [website](#).