



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Fas Ligand antibody [MFL3] (PE)

Cat No. GTX01516-08

| | |
|--------------------|------------------|
| Host | Armenian Hamster |
| Clonality | Monoclonal |
| Isotype | IgG |
| Application | FACS |
| Reactivity | Mouse |

Reference (1)

Package

100 µg

PRODUCT

Summary

The MFL3 monoclonal antibody reacts with mouse CD178, a 40 kD member type-II transmembrane protein and member of the TNF family of proteins. CD178 is also known as Fas ligand, FasL, Apo-1 ligand, and CD95 ligand. CD178 is expressed on activated T cells and natural killer (NK) cells and also tissue at immune privileged sites such as the eye and testis. FasL interacts with its receptor CD95 (Fas) to initiate apoptotic cell death and is thought to play a role in T cell development, immune response regulation, and cell-mediated cytotoxic responses. This MFL3 clone has been reported to block CD178/CD96 induced apoptosis.

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Dilution |
|--------------------|-----------------|
| FACS | Assay dependent |

Not tested in other applications.

Calculated MW 0|31 kDa. ([Note](#))


PROPERTIES

| | |
|----------------------|--|
| Form | Liquid |
| Buffer | 10 mM NaH ₂ PO ₄ (pH 7.2), 150 mM NaCl, 0.09% sodium azide, 0.1% gelatin |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light. |
| Concentration | 0.2 mg/ml (Please refer to the vial label for the specific concentration.) |
| Purification | Purified by affinity chromatography From tissue culture supernatant |
| Conjugation | Phycoerythrin (PE) |
| Note | For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

**GTX01516-08 FACS Image**

FACS analysis of mouse Fas Ligand transfected cells using GTX01516-08 Fas Ligand antibody [MFL3] (PE).

Solid line : primary antibody

Dashed line : isotype control

antibody amount : 0.25 μ g (5 μ l)



For full product information, images and publications, please visit our [website](#).