



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Dystrophin antibody [Dy10/12B2]

Cat No. GTX01869

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Application	IHC-Fr
Reactivity	Human

Package
500 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
IHC-Fr	Neat-1:20

Note : Freeze specimen tissue blocks in isopentane chilled in liquid nitrogen.

Not tested in other applications.

Calculated MW	427 kDa. (Note)
----------------------	-----------------------------------

Specificity/Sensitivity

This antibody reacts strongly with the amino terminal domain (between amino acids 321 and 494) of human dystrophin. Patient immunoreactivity indicates epitope is near exons 10 to 12. Epitope mapping suggests that sequences from amino acids 308 to 351 are involved in antibody binding. This region spans the junction of exons 9 and 10 and the epitope recognised may be part of a hinge region joining the amino domain to the central rod domain. No reactivity with DMD/BMD patients deleted for exons 10 to 12.

We do not recommend use of this product for Mouse,Rat,Rabbit,Dog,Hamster,Chicken,Pig.

PROPERTIES

Form	Liquid
Buffer	Tissue culture supernatant with sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	>4 µg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Fusion protein containing amino acids 67 to 713.
Purification	Unpurified
Conjugation	Unconjugated
Note	For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).