



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mucin 4 antibody [ZR201]

Cat. No. GTX03684

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Application	IHC-P
Reactivity	Human

Package
500 µl

PRODUCT

Summary

Mucin 4, a membrane-bound mucin, is the human homolog of the rat sialomucin complex (SMC). The Mucin 4 gene is the predominant mucin gene expressed in the normal urothelium and is also expressed in several normal tissues such as trachea, lung and testis. Dysregulation of mucin 4 results in high levels of expression in pancreatic tumors and tumor cell lines. Induction of mucin 4 in pancreatic carcinoma by all-trans-retinoic acid is mediated through the retinoic acid receptor- α signaling pathway. MUC-4 is a very specific (100%) and sensitive (90%) marker of lung adenocarcinomas and is negative for mesotheliomas. MUC4 is expressed in many human adenocarcinomas. It is specifically expressed in low grade fibromyxoid sarcoma. Reportedly, MUC-4 expression in invasive ductal carcinoma of the pancreas is an independent factor for poor prognosis.

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	Assay dependent

Not tested in other applications.

Calculated MW 232 kDa. ([Note](#))

PROPERTIES

Form	Liquid
Buffer	Tris-HCl buffer, 0.2% BSA
Preservative	0.09% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment (around aa 1730-1864) of human MUC4 prote
Purification	Purified IgG
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

Note For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).