



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UPK1B antibody [MSVA-734M] HistoMAX™

Cat. No. GTX04450

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Application	IHC-P
Reactivity	Human

Package

500 µl, 100 µl

PRODUCT

This antibody was validated on 76 different Normal Tissues by IHC-P.

Summary

[Go to Normal Tissue Gallery](#)
[Go to Cancer Tissue Gallery](#)

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100-1:200

Note : Manual staining : Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Tris-EDTA-based Target Retrieval Solution buffer.

This antibody has been validated by Autostainer including Agilent/Dako - Autostainer Link 48, Bond RX-Lecia, and Roche-Ventana Discovery ULTRA. The detailed steps by Autostainer please refer to the protocol.

Not tested in other applications.

Calculated MW 30 kDa. ([Note](#))

PROPERTIES

Form	Liquid
Buffer	PBS, 0.05% BSA (Please contact us for PBS only format)
Preservative	0.05% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment (around aa 109-229) of human Uroplakin 1B (UPK1B) protein
Purification	Protein A/G purified




For full product information, images and publications, please visit our [website](#).

Date 2024 / 06 / 27 Page 1 of 2

Conjugation Unconjugated


Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX04450 IHC-P Image

IHC-P analysis of human colon mucosa tissue section using GTX04450 UPK1B antibody [MSVA-734M] HistoMAX™.
UPK1B immunostaining is absent in all cells of the colon mucosa.



GTX04450 IHC-P Image

IHC-P analysis of human urinary bladder urothelium tissue section using GTX04450 UPK1B antibody [MSVA-734M] HistoMAX™.
A strong UPK1B immunostaining of all urothelial cells is seen in this urinary bladder sample.



GTX04450 IHC-P Image

IHC-P analysis of human urinary bladder muscle invasive urothelial carcinoma tissue section using GTX04450 UPK1B antibody [MSVA-734M] HistoMAX™.
Muscle invasive urothelial carcinoma of the urinary bladder showing a strong UPK1B immunostaining of tumor cells.



For full product information, images and publications, please visit our [website](#).