



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## EasyBlot anti Mouse IgG (HRP)

**Cat No. GTX221667-01**

<b>Application</b>	WB
<b>Reactivity</b>	Mouse

Reference ( 31 )

★★★★★ Review ( 1 )

Package

250 µl, 50 µl

## PRODUCT

## Summary

EasyBlot anti-Mouse IgG is an HRP-conjugated second step reagent that specifically reacts with the native, non-reduced form of mouse IgG and does not bind the reduced, denatured form. Using Easyblot decreases the interference caused by the heavy (~50 kDa) and light chains (~25 kDa) of the IgG used for immunoprecipitation (IP). EasyBlot is simple to use and results in Western Blot detection of only the target band in your IP experiments. Exercise caution when using Protein A/G for IP with EasyBlot. Protein A/G contaminants bind to mouse IgG with high affinity and may cause contaminating bands with a molecular weight similar to IgG heavy chain or light chain.

To minimize background from Protein A/G contamination, we strongly recommend the use of EasyBlot Kit anti-Mouse IgG (Optimized for Prot A/G) (GTXGTX225857-01) or EasyBlocker (GTX425858) to minimize protein A/G contamination or you can use anti-Mouse IgG Agarose beads (GTX27018) instead of Protein A/G.

## APPLICATION

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

## Suggested dilution

Suggested dilution	Dilution
WB	1:1000

**Note : Use 2-5 µg of the IP antibody; do not exceed 5 µg. Completely reduce all IP samples by 100mM DTT (fresh prepared). For further detail, please refer to the protocol.**

Not tested in other applications.

## PROPERTIES

**Form** Liquid

**Buffer** 0.05M Tris (pH7.4), 0.15M NaCl, 1%BSA, 40% Glycerol

**Preservative** 0.025% ProClin 300

**Storage** Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

**Concentration** 0.03 mg/ml (Please refer to the vial label for the specific concentration.)


**Purification** Purified by antigen-affinity chromatography.

**Conjugation** Horseradish peroxidase(HRP)


**Note** For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

**DATA IMAGES**

**GTX221667-01 WB Image**

Western Blot analysis of native mouse IgG (left) or denatured mouse IgG (right) using EasyBlot anti Mouse IgG (HRP) antibody.


**GTX221667-01 WB Image**

EasyBlot anti-Mouse IgG (HRP) (GTX221667-01) was used as a secondary reagent for immunoprecipitation analysis. 293T lysates were immunoprecipitated using 5 ug of normal mouse IgG or ELP3 antibody [GT11811] (GTX631395)



For full product information, images and publications, please visit our [website](#).