



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## C1q antibody

Cat. No. GTX28779

<b>Host</b>	Sheep
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Application</b>	Double immunodiffusion, RID
<b>Reactivity</b>	Human

Package  
1 ml

## APPLICATION

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
Double immunodiffusion	Assay dependent
RID	Neat-1:2

**Note : 0.1 µl antiserum / cm<sup>2</sup> gel vs. 20 µl fresh plasma Neat-1:2**

Not tested in other applications.

## Product Note

Gives a single arc when tested by IEP against fresh human plasma. Identity has been confirmed by double diffusion (Ouchterlony) against fresh human plasma and a known anti-human C1q. C1q is a sub-component of the first complement component C1, which is formed by the interaction of C1q with the tetramer C1s-C1r-C1r-C1s. C1q is a member of a family of soluble collagen-like proteins and is composed of three basic subunits termed A, B and C (27,550, 25,200, 23,800 Da- respectively) which associate in the proportion 6 A-B subunits (52,750 Da) and 3 C-C subunits (47,600 Da), to form whole C1q (459,300 Da). C1q binds to IgG and IgM immune complexes, this activates C1r and C1s, which are also bound to C1q and so initiates the classical pathway of complement activation. The normal plasma concentration of C1q is 100-180 mg/L.

## PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	Glycine buffered Saline pH7.4, 0.1% EACA, 0.01% Benzamidine, 1mM EDTA
<b>Preservative</b>	0.099% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
<b>Concentration</b>	5.07 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Human C1q, > 95% pure (SDS-PAGE). Prepared from human plasma.
<b>Purification</b>	IgG fraction From serum
<b>Conjugation</b>	Unconjugated



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).