

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PER2 antibody [HL2775]

Cat. No. GTX639640

| Host | Rabbit |
|-------------|------------|
| Clonality | Monoclonal |
| lsotype | lgG |
| Application | WB |
| Reactivity | Human |

Package

100 µl, 25 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|-----------------------------------|----------------------|
| WB | 1:500-1:3000 |
| Not tested in other applications. | |

Calculated MW


137 kDa. (<u>Note</u>)

| PROPERTIES | | |
|---------------|--|--|
| Form | Liquid | |
| Buffer | PBS | |
| Preservative | No preservative | |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. | |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) | |
| Immunogen | Recombinant fragment of human PER2. | |
| Purification | Affinity purified by Protein A. | |
| Conjugation | Unconjugated | |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. | |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. | |




For full product information, images and publications, please visit our <u>website</u>.

DATA IMAGES




GTX639640 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with PER2 antibody [HL2775] (GTX639640) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX639640 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with PER2 antibody [HL2775] (GTX639640) diluted at 1:100000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX639640 WB Image

Treated 293T whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with PER2 antibody [HL2775] (GTX639640) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 06 / 27 Page 2 of 2