



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Anti-TCR Beta Antibody (H57-597)

Description

Anti-TCR Beta Antibody (H57-597)

Product Benefits:

ichorbio's Anti-TCR Beta Antibody (H57-597) can be used for in vivo use as it has low endotoxin, no BSA or preservatives.

Target:

TCR Beta

Clone:

H57-597

Size:

ichorbio's Anti-Mouse TCR Beta In Vivo Antibody – Low Endotoxin (H57-597) is available in the following sizes:

5mg, 10mg, 20mg, 50mg

Isotype:

Armenian Hamster IgG

Other names:

TCR-? chain, TCR-?, ?-TCR

Host:

Armenian Hamster

Species Reactivity:

Mouse

Specificity:

Clone H57-597 recognizes an epitope on the constant region of mouse TCR β . Clone H57-597 does not cross-react with α T cells.

Purification Method:

This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.

Antigen Distribution:

TCR β is expressed on α thymocytes, peripheral T cells, and NKT cells.

Background:

TCR β antibody, clone H57-597, recognizes the beta chain of the T cell receptor (TCR β), expressed on α thymocytes and T lymphocytes, as well as natural killer T (NKT) cells. TCR β is a transmembrane protein that consists of an amino-terminal variable region, a constant region, and a stalk segment covalently linked to the TCR alpha chain (TCR α) via a disulfide bond. TCR β does not contain signaling domains and is instead noncovalently associated with the CD3 complex (TCR-CD3), essential for TCR signaling. TCR β recognizes antigens bound to major histocompatibility complex (MHC) molecules, resulting in phosphorylation of CD3 ITAMs and subsequent T cell activation, proliferation, and survival.

Immunogen:

Affinity purified TCR from mouse DO-11.10 cells

Concentration:

5mg/ml

Formulation:

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 – 7.4 with no carrier protein, potassium, calcium or preservatives added.

Purity:

\geq 95% monomer by analytical SEC, \geq 95% by SDS Page

Endotoxin:

? 1.0 EU/mg as determined by the LAL method

Storage:

This antibody is stable for at least one week when stored sterile at 2-8°C. For long term storage aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Applications:

Flow Cytometry, Costimulatory, Depletion, Immunoprecipitation, IHC (Frozen), IHC (Paraffin), CODEX.

Application notes:

The suggested concentration for this TCR ? chain (clone H57-597) antibody for staining cells in flow cytometry is ? 0.25 ?g per 10⁶ cells in a volume of 100 ?l. Titration of the reagent is recommended for optimal performance for each application.

Isotype Control:

[Armenian Hamster IgG Isotype Control for In Vivo – Low Endotoxin \[PIP\] \(ICH2251\)](#)