



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse Anti-Mouse PD-1 Antibody (RMP1-14) D265A

Description

Mouse Anti-Mouse PD-1 Antibody (RMP1-14) D265A

Product Benefits:

ichorbio's mouse anti-mouse PD-1 antibody (RMP1-14) D265A is manufactured in a cGMP compliant facility. Click [here](#) for ichorbio's RMP1-14 antibodies and [here](#) to view ichorbio's complete list of anti-PD-1 antibodies and biosimilars. ichorbio: the best antibodies for *in vivo* research.

Size:

ICH1182D265A mouse anti-mouse RMP1-14 is available in the following sizes:
5mg, 10mg, 20mg and 50mg

Target:

PD-1

Clone:

RMP1-14

Isotype:

Mouse IgG1

Other Names:

Programmed cell death protein 1, Pcd1, CD279

Uniprot:

Q02242

Host:

CHO

Species Reactivity:

Mouse

Specificity:

Mouse anti-mouse PD-1 antibody (RMP1-14) recognizes an epitope on Mouse PD-1. ICH1182D265A contains a mutation at D265A to reduce its effector function.

Purification Method:

This recombinant antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.

Antigen Distribution:

Subset of double negative thymocytes, activated T and B cells

Background:

CD279, also known as programmed death-1 (PD-1) is a 50-55 kD immunoglobulin superfamily member. PD-1 is expressed on a subset of CD4-CD8- thymocytes, and on activated T and B cells. The PD-1 ligands, PD-L1 (also known as B7-H1) and PD-L2 (B7-DC), are members of the B7 immunoglobulin superfamily. This RMP1-14 antibody has been reported to block the binding of PD-1 to its ligands (B7-H1 and B7-DC) and to inhibit T cell proliferation and cytokine production.

Immunogen:

Mouse PD-1 transfected BHK cells

Concentration:

1.0 – 5.0 mg/ml

Formulation:

0.01 M phosphate buffered saline (PBS) pH 7.2, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and Azide free.

Purity:

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

Endotoxin:

? 1.0 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

Storage:

This antibody is stable for at least 4 weeks when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at – 20°C or -80°C. Avoid repeated freeze thaw cycles.

Applications:

Western Blot, Blocking, Flow Cytometry, Functional Assays

Use:

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

Isotype Control:

[Mouse IgG1 Isotype Control for In Vivo – Low Endotoxin \(HKSP84\) \[ICH2248\]](#)

Original Antibody:

[Anti-PD-1 In Vivo Antibody – Low Endotoxin \[RMP1-14\]](#)

Antibodies against the same target:

[Anti-PD-1 In Vivo Antibody – Low Endotoxin \[29F.1A12\] \(ICH1091\)](#), [Anti-PD-1 In Vivo Antibody – Ultra Low Endotoxin \[29F.1A12\] \(ICH1091UL\)](#)