



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Name: Anti-Mouse 4-1BBL In Vivo Antibody - Low Endotoxin (TKS-1) [ICH1184]

Sku: ICH1184

Catagories: [anti-mouse](#), [Blocking](#), [Functional Assays](#), [Low Endotoxin](#), [Ultra Low Endotoxin](#), [Western Blot](#)



Product price: \$375.00 – \$2,950.00

Product Details

Anti-Mouse 4-1BBL In Vivo Antibody - Low Endotoxin (TKS-1)

Bio X Cell:

ICH1184 is [up to 30% cheaper](#) for academia & non-profits and [up to 55% cheaper](#) for industry than the equivalent product BE0110 from Bio X Cell.

Product Benefits:

ichorbio's Anti-Mouse 4-1BBL In Vivo Antibody - Low Endotoxin (TKS-1) [ICH1184] is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from our [competitors](#) at less than 1.0 EU/mg. If ichorbio's low endotoxin antibodies are not low enough we also offer ultra low endotoxin antibodies which have even less endotoxin (<0.75EU/mg) at an even higher purity (98% versus 95%). ichorbio: the best antibodies for *in vivo* research.

Target:

4-1BBL

Clone:

TKS-1

Size:

ichorbio's TKS-1 *in vivo* antibody is available in the following bulk sizes: 5mg, 25mg, 50mg and 100mg. ichorbio regularly manufactures multi-gram amounts of our anti-4-1BBL TKS-1 clone - please contact us for pricing.

Isotype:

IgG2a kappa

Other Names:

CD137L, TNFSF9, 4-1BB Ligand, TKS-1

Host:

Rat

Species Reactivity:

Mouse

Antigen Distribution:

4-1BBL is expressed by activated B cells, macrophages, and dendritic cells.

Background:

4-1BBL (CD137L) and 4-1BB (CD137) are a receptor ligand pair belonging to the tumor necrosis factor (TNF) receptor/TNF superfamily. 4-1BBL is a type II transmembrane protein expressed by activated B cells, macrophages, and dendritic cells. Members of the TNF-TNF receptor superfamily play critical roles in regulating cellular activation, differentiation, and apoptosis and the 4-1BBL-4-1BB interaction is important in cellular immune responses. 4-1BBL-4-1BB interaction provides a co-stimulatory signal to T cells, resulting in increased proliferation and cytokine production. A TKS-1-producing hybridoma was generated by immunizing a SD rat with 4-1BBL-transfected NRK cells (rat IgG2a,?). Splenocytes were fused with P3U1 cells and hypoxanthine-aminopterin-thymidine selection was performed. The monoclonal antibody was identified by its strong reactivity with 4-1BBL-transfected L5178Y cells, cloned by limiting diffusion, purified from ascites, and found to bind to 4-1BBL/NRK and 4-1BBL/P815 cells. Pre-incubation with TKS-1 blocks 4-1BB-Ig binding to 4-1BBL/L5178Y cells, indicating TKS-1 is specific to mouse 4-1BBL and can interrupt the interaction between 4-1BBL and 4-1BB. Additionally, TKS-1 can block anti-CD3-stimulated T cell proliferation co-stimulated by 4-1BBL /P815 cells. The monoclonal antibodies 19H3 and TKS-1 bind to different sites on murine 4-1BBL, with TKS-1 binding to a site that overlaps with the receptor binding site. Additionally, TKS-1 and 4-1BB bind to a similar site, making TKS-1 especially useful for blocking studies.

Concentration:

1.0 - 5.0 mg/ml

Formulation:

0.01 M phosphate buffered saline (PBS) pH 7.2, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and Azide free.

Purity:

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

Endotoxin:

<1.0 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

Aggregation:

Aggregation level ? 5%

Aggregation level ? 1%

IMPACT Pathogen Test:

We use the IMPACT test generated by IDEXX Laboratories to guarantee our Ultra Low Endotoxin antibodies are pathogen free. Our mouse antibodies are tested for: Mycoplasma spp., Mycoplasma pulmonis, Sendai virus, Mouse hepatitis virus, Pneumonia virus of mice, Minute virus of mice, Mouse parvovirus (MPV1-5), Theiler's murine encephalomyelitis virus, Murine norovirus, Reovirus 3, Mouse rotavirus, Ectromelia virus, Lymphocytic choriomeningitis virus, Polyoma virus, Lactate dehydrogenase-elevating virus, Mouse adenovirus (MAD1, MAD2), Mouse cytomegalovirus, K virus, Mouse thymic virus, Hantaan virus, Corynebacterium bovis and Corynebacterium spp. (HAC2)

Storage:

Anti-4-1BBL In Vivo Antibody - Low Endotoxin (TKS-1) is stable for at least one week when stored sterile at 2-8°C. For long term storage aseptically aliquot in working volumes without diluting and store at -20°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

Applications:

Functional Assays, Blocking, Western Blot

Application Notes:

Each investigator should determine their own optimal working dilution for specific applications..

Use:

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

Isotype Control:

[Rat IgG2a In Vivo Isotype Control – Low Endotoxin \(1-1\) \[ICH2244\]](#)

Additional information:

Size: [5mg](#), [25mg](#), [50mg](#), [100mg](#)

Endotoxin Level: Low, Ultra Low