



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti- Human CD55 (JS11)

Fluorochrome	Reference	Size
FITC	55F-100T	100 test

PRODUCT DESCRIPTION

Clone: JS11

Isotype: IgG1

Tested application: flow cytometry

Immunogen: The anti-CD55 monoclonal antibody derives from Human Peripheral Blood Mononuclear Cells and Tonsil cells.

Species reactivity: Human

Storage instruction: store in the dark at 2-8 °C

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃).

Recommended usage: Immunostep's CD55, clone JS11 is a monoclonal antibody intended for the identification and enumeration of complement decay-accelerating factor protein expressed on hematopoietic cells including erythrocytes and many non-hematopoietic cells using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using $\leq 1 \mu\text{g}/10^6$ cells.

Presentation: liquid

Source: Supernatant proceeding from an *in vitro* cell culture of a cell hybridoma.

Purification: Affinity chromatography.

ANTIGEN DETAILS

Large description: CD55 clone JS11 reacts with DAF protein a glycosylphosphatidylinositol (GPI)-anchored single chain glycoprotein also known as decay-accelerating factor.

It is expressed on hematopoietic cells including erythrocytes and many non-hematopoietic cells. This protein recognizes C4b and C3b fragments that condense with cell-surface hydroxyl or amino groups when nascent C4b and C3b are locally generated during C4 and c3 activation. Interaction of DAF with cell-associated C4b and C3b polypeptides interferes with their ability to catalyze the conversion of C2 and factor B to enzymatically active C2a and Bb and thereby prevents the formation of C4b2a and C3bBb, the amplification convertases of the complement cascade. CD55 has been reported to reduce the efficiency of NK cell lysis and induce signal transduction in T cells. CD55 has also been shown to interact with CD97 and bind to Coxsackie and Echovirus.⁽¹⁻³⁾

Other Names: Complement decay-accelerating factor, DAF

Gene ID: 1604

Molecular weight: 41,4 kDa

Please, refer to www.immunostep.com technical support for more information.

WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties. Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

REFERENCES

1. Nowicki B, Nowicki S. DAF as a therapeutic target for steroid hormones: implications for host-pathogen interactions. *Adv Exp Med Biol*;735:83-96.
2. Caras IW, Davitz MA, Rhee L, Weddell G, Martin DW, Jr., Nussenzweig V. Cloning of decay-accelerating factor suggests novel use of splicing to generate two proteins. *Nature*1987 Feb 5-11;325(6104):545-9.
3. Osuka F, Endo Y, Higuchi M, Suzuki H, Shio Y, Fujii K, et al. Molecular cloning and characterization of novel splicing variants of human decay-accelerating factor. *Genomics*2006 Sep;88(3):316-22.

MANUFACTURED BY



Immunostep S.L.
Avda. Universidad de Coimbra, s/n
Cancer Research Center (CIC)
Campus Miguel de Unamuno
37007 Salamanca (Spain)
Tel. (+34) 923 294 827
www.immunostep.com