



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Anti- Human CD38 (GR7-A4)(LD38)

Fluorochrome	Reference	Size
Pure	38PUI-OIMG	100 test
FITC	38F1-I00T	100 test
PE	38PE1-I00T	100 test
PerCP	38PP1-I00T	100 test
APC	38AI-I00T	100 test
DY634	38DY1-I00T	100 test
Biotin	38B1-OIMG	100 test
CF-Blue	38CFB1-I00T	100 test
OC515	38OC1-I00T	100 test
APC-C750	38AC7501-I00T	100 test

PRODUCT DESCRIPTION

Clone: GR7-A4

Isotype: IgG1

Tested application: flow cytometry

Immunogen: The anti-CD38 monoclonal antibody derives from cells of human acute lymphoblastic B leukaemia (B-ALL).

Species reactivity: Human

Storage instruction: store in the dark at 2-8 °C

Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₃).

Recommended usage: Immunostep's CD38, clone GR7-A4, is a monoclonal antibody intended for the identification and enumeration of plasma cells using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using $\leq 1 \mu\text{g}/10^6$ cells.

Presentation: liquid

Source: Supernatant proceeding from an *in vitro* cell culture of a cell hybridoma.

Purification: Affinity chromatography.

ANTIGEN DETAILS

Large description: The monoclonal antibody is directed against the T10-antigen, which is strongly expressed plasma cell. The CD38 antigen is expressed at variable levels on the majority of hemopoietic cells, prevalently during early differentiation and activation.

CD38 synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.⁽¹⁻⁵⁾

Other Names: ADP-ribosyl cyclase/cyclic, ADP-ribosyl cyclase 1, ADPRC 1, Cyclic ADP-ribose hydrolase 1, cADPr hydrolase 1, T10.

Gene ID: 952

Molecular weight: 45 kDa

Please, refer to www.immunostep.com technical support for more information.

WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties. Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

REFERENCES

1. Ferrero E, Malavasi F. The metamorphosis of a molecule: from soluble enzyme to the leukocyte receptor CD38. *J Leukoc Biol* 1999 Feb;65(2):151-61.
2. Lund F, Solvason N, Grimaldi JC, Parkhouse RM, Howard M. Murine CD38: an immunoregulatory ectoenzyme. *Immunol Today* 1995 Oct;16(10):469-73.
3. Malavasi F, Funaro A, Roggero S, Horenstein A, Calosso L, Mehta K. Human CD38: a glycoprotein in search of a function. *Immunol Today* 1994 Mar;15(3):95-7.
4. Mehta K, Shahid U, Malavasi F. Human CD38, a cell-surface protein with multiple functions. *FASEB J* 1996 Oct;10(12):1408-17.
5. Shubinsky G, Schlesinger M. The CD38 lymphocyte differentiation marker: new insight into its ectoenzymatic activity and its role as a signal transducer. *Immunity* 1997 Sep;7(3):315-24.

MANUFACTURED BY



Immunostep S.L

Avda. Universidad de Coimbra, s/n
 Cancer Research Center (CIC)
 Campus Miguel de Unamuno
 37007 Salamanca (Spain)
 Tel. (+34) 923 294 827
www.immunostep.com