



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HEPES sodium

Cat. No.:	HY-108535
CAS No.:	75277-39-3
Molecular Formula:	C ₈ H ₁₇ N ₂ NaO ₄ S
Molecular Weight:	260.29
Target:	Biochemical Assay Reagents
Pathway:	Others
Storage:	Store at room temperature



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (96.05 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	3.8419 mL	19.2093 mL	38.4187 mL
				5 mM	0.7684 mL	3.8419 mL	7.6837 mL
				10 mM	0.3842 mL	1.9209 mL	3.8419 mL
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (9.60 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (9.60 mM); Clear solution						
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (9.60 mM); Clear solution						

BIOLOGICAL ACTIVITY

Description	HEPES sodium, a nonvolatile zwitterionic chemical buffering agent, is broadly applied in cell culture. HEPES sodium is effective at pH 6.8 to 8.2. HEPES sodium is also a potent inducer of lysosome biogenesis ^{[1][2][3]} .
In Vitro	HEPES maintains superhydrophilicity of titanium for at least 3 months and resulted in a continuous retention of bioactivity and osteoconductivity ^[1] . HEPES drives lysosome biogenesis, affects MiT/TFE cytoplasmic-nuclear distribution, disrupts global cellular transcriptional profiles, resulting the activation of a MiT/TFE-dependent lysosomal-autophagic gene network in cultured RAW264.7 cells ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Int Immunopharmacol. 2023 May 12;120:110292.
- Int Immunopharmacol. September 2022, 108953.
- Andrology. 2024 May 22.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Suzuki T, et al. Nonvolatile buffer coating of titanium to prevent its biological aging and for drug delivery. Biomaterials. 2010;31(18):4818-4828.
- [2]. Sledź P, et al. An experimental charge density of HEPES. Acta Crystallogr B. 2010;66(Pt 4):482-492.
- [3]. <https://pubmed.ncbi.nlm.nih.gov/20631430/>
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA