



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## BI-3802

Cat. No.:	HY-108705		
CAS No.:	2166387-65-9		
Molecular Formula:	C <sub>24</sub> H <sub>29</sub> ClN <sub>6</sub> O <sub>3</sub>		
Molecular Weight:	484.98		
Target:	Bcl-2 Family; Molecular Glues		
Pathway:	Apoptosis; PROTAC		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 6.67 mg/mL (13.75 mM; ultrasonic and warming and heat to 60°C)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	2.0619 mL	10.3097 mL	20.6194 mL
5 mM	0.4124 mL	2.0619 mL	4.1239 mL
10 mM	0.2062 mL	1.0310 mL	2.0619 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

BI-3802 is a highly potent BCL6 degrader and inhibits the Bric-à-brac (BTB) domain of BCL6 with an IC<sub>50</sub> of ≤3 nM. BI-3802 induces the polymerization of BCL6 and promotes BCL6 degradation depended on E3 ligase SIAH1. BI-3802 has antitumor activity<sup>[1][2]</sup>.

#### IC<sub>50</sub> & Target

IC<sub>50</sub>: ≤ 3 nM (BCL6 BTB)<sup>[1]</sup>

#### In Vitro

BI-3802 shows an IC<sub>50</sub> of 43 nM for the cellular BCL6<sup>[1]</sup>.  
 BI-3802 increases the interaction between BCL6 and SIAH1 (EC<sub>50</sub> = 64 nM)<sup>[2]</sup>.  
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Kerres N, et al. Chemically Induced Degradation of the Oncogenic Transcription Factor BCL6. Cell Rep. 2017 Sep 19;20(12):2860-2875.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA