



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rp-8-pCPT-cGMPS sodium

Cat. No.:	HY-110221
CAS No.:	208445-07-2
Molecular Formula:	C ₁₆ H ₁₄ ClN ₅ NaO ₆ PS ₂
Molecular Weight:	525.86
Target:	PKG
Pathway:	Stem Cell/Wnt
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Rp-8-pCPT-cGMPS sodium is the sodium salt form of Rp-8-pCPT-cGMPS. Rp-8-pCPT-cGMPS is an inhibitor for cGMP-dependent protein kinase (cGK). Rp-8-pCPT-cGMPS sodium is an agonist for cyclic nucleotide-gated (CNG) channels in a voltage-dependent manner ^{[1][2]} .	
In Vivo	Rp-8-pCPT-cGMPS sodium (5 nM, intrathecally, once a day for 3 days) inhibits the cGMP-cGKI pathway, suppresses tumor cell implantation (TCI)-induced thermal hyperalgesia and mechanical allodynia in Sprague-Dawley rats model ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	Bone cancer pain induced by tumor cell implantation (TCI) in Sprague-Dawley rats ^[1] .
	Dosage:	0.5 mmol/L, 20 μL
	Administration:	intrathecally, once a day for 3 days
	Result:	Delayed and suppressed TCI-induced thermal hyperalgesia and mechanical allodynia.

REFERENCES

[1]. Liu S, et al., cGMP and cGMP-dependent protein kinase I pathway in dorsal root ganglia contributes to bone cancer pain in rats. *Spine (Phila Pa 1976)*. 2014 Sep 1;39(19):1533-41.

[2]. Wei JY, et al., Substituted cGMP analogs can act as selective agonists of the rod photoreceptor cGMP-gated cation channel. *J Mol Neurosci*. 1998 Feb;10(1):53-64.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA