



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

## UDP-Galactose disodium

<b>Cat. No.:</b>	HY-114364
<b>CAS No.:</b>	137868-52-1
<b>Molecular Formula:</b>	C <sub>15</sub> H <sub>22</sub> N <sub>2</sub> Na <sub>2</sub> O <sub>17</sub> P <sub>2</sub>
<b>Molecular Weight:</b>	610.27
<b>Target:</b>	P2Y Receptor; Endogenous Metabolite
<b>Pathway:</b>	GPCR/G Protein; Metabolic Enzyme/Protease
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 250 mg/mL (409.65 mM; Need ultrasonic)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	1.6386 mL	8.1931 mL	16.3862 mL
5 mM	0.3277 mL	1.6386 mL	3.2772 mL
10 mM	0.1639 mL	0.8193 mL	1.6386 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

UDP-Galactose disodium is a monosaccharide and a P2Y14 receptor agonist with an EC<sub>50</sub> value of 0.67 μM. UDP-Galactose disodium is a substrate for the transferase beta-1, 4 galactosyltransferase V (B4GALT5)<sup>[1][2]</sup>.

#### IC<sub>50</sub> & Target

P2Y14 Receptor  
0.67 μM (EC50)

### REFERENCES

- [1]. Ko H, et al. Structure-activity relationship of uridine 5'-diphosphoglucose analogues as agonists of the human P2Y14 receptor. *J Med Chem.* 2007 May 3;50(9):2030-9.
- [2]. Drabavicius G, Daelemans D. Intermedilysin cytolytic activity depends on heparan sulfates and membrane composition. *PLoS Genet.* 2021 Feb 12;17(2):e1009387.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA