



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## 2-Iminobiotin

Cat. No.:	HY-118700		
CAS No.:	13395-35-2		
Molecular Formula:	C <sub>10</sub> H <sub>17</sub> N <sub>3</sub> O <sub>2</sub> S		
Molecular Weight:	243.33		
Target:	NO Synthase		
Pathway:	Immunology/Inflammation		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : 12.5 mg/mL (51.37 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	4.1096 mL	20.5482 mL	41.0965 mL
	5 mM	0.8219 mL	4.1096 mL	8.2193 mL
	10 mM	0.4110 mL	2.0548 mL	4.1096 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

2-Iminobiotin (Guanidinobiotin) is a biotin (vitamin H or B7) analog. 2-Iminobiotin is a reversible nitric oxide synthases inhibitor with K<sub>s</sub> of 21.8 and 37.5 μM for murine iNOS and rat n-cNOS, respectively<sup>[1]</sup>. 2-Iminobiotin superimposes on hypothermia protects human neuronal cells from hypoxia-induced cell damage<sup>[2]</sup>.

#### IC<sub>50</sub> & Target

iNOS

#### In Vitro

Application of low concentrations of 2-Iminobiotin (2-IB; 10 ng/mL and 30 ng/mL) superimposed on hypothermia abrogates the hypoxia-induced lactate dehydrogenase (LDH) increase, resulting in LDH levels that were not different from the normoxia control<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Sup SJ, et al. 2-Iminobiotin is an inhibitor of nitric oxide synthases. *Biochem Biophys Res Commun.* 1994 Oct 28;204(2):962-8.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA