



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## IWR-1

<b>Cat. No.:</b>	HY-12238		
<b>CAS No.:</b>	1127442-82-3		
<b>Molecular Formula:</b>	C <sub>25</sub> H <sub>19</sub> N <sub>3</sub> O <sub>3</sub>		
<b>Molecular Weight:</b>	409.44		
<b>Target:</b>	Wnt; Organoid		
<b>Pathway:</b>	Stem Cell/Wnt		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 50 mg/mL (122.12 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	2.4424 mL	12.2118 mL	24.4236 mL
		5 mM	0.4885 mL	2.4424 mL	4.8847 mL
10 mM		0.2442 mL	1.2212 mL	2.4424 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	<ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: ≥ 2.5 mg/mL (6.11 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.11 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: ≥ 2.5 mg/mL (6.11 mM); Clear solution</li> </ol>				

### BIOLOGICAL ACTIVITY

<b>Description</b>	IWR-1 is a tankyrase inhibitor which inhibits Wnt/β-catenin signaling pathway.
<b>IC<sub>50</sub> &amp; Target</b>	IC <sub>50</sub> : 180 nM (Wnt)
<b>In Vitro</b>	Both IWR-1 and XAV939 act as reversible Wnt pathway inhibitors and exhibit similar pharmacological effects in vitro. IWR-1 exerts its effect via interaction with Axin, while XAV939 binds TNKS directly <sup>[1]</sup> . IWR-1 (10 μM) induces stabilization of β-catenin disruption complex. IWR-1 (10 μM) is added to the medium together with MIF, the size of cell colonies is extremely

---

decreased, and that indicates the promoting effect of MIF on NSPC proliferation is inhibited by IWR-1 in any MIF concentration group. 2, 5 and 10  $\mu$ M of IWR-1 significantly inhibits the proliferation of NSPC dose-dependently. IWR-1 inhibits the promoting effect of MIF on NSPC differentiation to neuron lineage<sup>[2]</sup>. IWR-1 treatment in the presence of maximal stimulatory dose of FSH (0.5 ng/mL) results in a dose dependent inhibition of the stimulatory effect of FSH with > 75% inhibition observed at the maximal inhibitory dose of IWR-1 (1  $\mu$ M). IWR-1 treatment partially reverses the FSH-induced inhibition of granulosa cell CARTPT mRNA expression<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## CUSTOMER VALIDATION

- Adv Funct Mater. 2023 Dec 22.
- Nano Today. 21 September 2022.
- Nat Commun. 2024 May 23;15(1):4393.
- Sci Total Environ. 2022 Feb 25;809:152102.
- Chemosphere. 2023 Sep 27:140283.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Lu J, et al. Structure-activity relationship studies of small-molecule inhibitors of Wnt response. *Bioorg Med Chem Lett*. 2009 Jul 15;19(14):3825-7.

[2]. Zhang X, et al. Macrophage migration inhibitory factor promotes proliferation and neuronal differentiation of neural stem/precursor cells through Wnt/ $\beta$ -catenin signal pathway. *Int J Biol Sci*. 2013 Nov 28;9(10):1108-20.

[3]. Gupta PS, et al. Regulation and Regulatory Role of WNT Signaling in Potentiating FSH Action during Bovine Dominant Follicle Selection. *PLoS One*. 2014 Jun 17;9(6):e100201.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA