



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AE-ITU dihydrobromide

Cat. No.:	HY-131929
CAS No.:	56-10-0
Molecular Formula:	C ₃ H ₁₁ Br ₂ N ₃ S
Molecular Weight:	281.01
Target:	NO Synthase
Pathway:	Immunology/Inflammation
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	AE-ITU dihydrobromide is the dihydrobromide form of AE-ITU. AE-ITU dihydrobromide is a selective inhibitor for inducible NO synthase (iNOS), and attenuates the liver dysfunction caused by endotoxaemia in rats ^[1] .								
IC₅₀ & Target	iNOS								
In Vivo	<p>AE-ITU dihydrobromide (1 mg/kg/h, i.v., for 1 hour) attenuates the delayed hypotension and vascular hyporeactivity, ameliorates liver dysfunction in LPS (HY-D1056)-administrated Wistar rats^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>LPS (HY-D1056)-administrated Wistar rats^[1]</td> </tr> <tr> <td>Dosage:</td> <td>1 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>i.v., 1 mg/kg/h for 1 hour</td> </tr> <tr> <td>Result:</td> <td>Reduced the serum levels of bilirubin, serum nitrite, GOT, GPT and γGT caused by LPS, inhibited iNOS in lung and liver homogenates.</td> </tr> </table>	Animal Model:	LPS (HY-D1056)-administrated Wistar rats ^[1]	Dosage:	1 mg/kg	Administration:	i.v., 1 mg/kg/h for 1 hour	Result:	Reduced the serum levels of bilirubin, serum nitrite, GOT, GPT and γGT caused by LPS, inhibited iNOS in lung and liver homogenates.
Animal Model:	LPS (HY-D1056)-administrated Wistar rats ^[1]								
Dosage:	1 mg/kg								
Administration:	i.v., 1 mg/kg/h for 1 hour								
Result:	Reduced the serum levels of bilirubin, serum nitrite, GOT, GPT and γGT caused by LPS, inhibited iNOS in lung and liver homogenates.								

REFERENCES

[1]. Thiemeermann C, et al., The multiple organ dysfunction syndrome caused by endotoxin in the rat: attenuation of liver dysfunction by inhibitors of nitric oxide synthase. Br J Pharmacol. 1995 Dec;116(7):2845-51.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA