



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## 3,4-Dehydro Cilostazol-d<sub>11</sub>

Cat. No.:	HY-135910S
CAS No.:	1073608-13-5
Molecular Formula:	C <sub>20</sub> H <sub>14</sub> D <sub>11</sub> N <sub>5</sub> O <sub>2</sub>
Molecular Weight:	378.51
Target:	Drug Metabolite; Isotope-Labeled Compounds
Pathway:	Metabolic Enzyme/Protease; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	3,4-Dehydro Cilostazol-d <sub>11</sub> is deuterated labeled 3,4-Dehydro Cilostazol (HY-135910). 3,4-Dehydro Cilostazol (OPC-13015) is an active metabolite of Cilostazol (CLZ; HY-17464). 3,4-Dehydro Cilostazol is used for pharmacokinetic study <sup>[1]</sup> .
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
<b>In Vivo</b>	3,4-Dehydro Cilostazol (OPC-13015; Oral; 1 mg/kg GLZ and 10 mg/kg CLZ) has a T <sub>1/2</sub> of 3.94 hours, a C <sub>max</sub> of 1.39 μM and an AUC <sub>0-24</sub> of 4.69 μg <sup>2</sup> h/ml. The plasma concentration time profiles of GLZ, CLZ & its active metabolite DCLZ are traceable up to 24 h, 12 h and 12 h respectively by oral administration at 1 mg/kg dose of GLZ and 10 mg/kg CLZ <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. T R S Satheeshmanikandan, et al. Liquid Chromatography - Tandem Mass Spectrometry for the Simultaneous Quantitation of Glipizide, Cilostazol and Its Active Metabolite 3, 4-dehydro-cilostazol in Rat Plasma: Application for a Pharmacokinetic Study. *Arzneimittelforschung*. 2012 Sep;62(9):425-32.

[2]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019 Feb;53(2):211-216.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA