



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IWP-2 (GMP)

Cat. No.:	HY-13912G
CAS No.:	686770-61-6
Molecular Formula:	C ₂₂ H ₁₈ N ₄ O ₂ S ₃
Molecular Weight:	466.6
Target:	Wnt
Pathway:	Stem Cell/Wnt
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	IWP-2 (GMP) is IWP-2 (HY-13912) produced by using GMP guidelines. GMP small molecules work appropriately as an auxiliary reagent for cell therapy manufacture. IWP-2 is an inhibitor of Wnt processing and secretion with an IC ₅₀ of 27 nM. IWP-2 targets the membrane-bound O-acyltransferase porcupine (Porcn) and blocks Wnt ligand palmitoylation ^[1] .								
IC₅₀ & Target	IC ₅₀ : 27 nM (Wnt) ^[1]								
In Vitro	<p>IWP-2 (GMP) (2 μM, in the first 4 days of Stage IV induction medium) reprograms human somatic cells to pluripotent stem cells^[1].</p> <p>IWP-2 (GMP) (5 μM, day 3 to 5) induces cardiac differentiation of hiPSCs^[2].</p> <p>IWP-2 (GMP) (5 μM, day 1 to 3) increasing the expression of cardiac progenitors and cardiac genes (MYL2, TNNT3, and TNNT2) in hiPSCs^[3].</p> <p>IWP-2 (GMP) (5 μM, day 5-7) together with PIP-S2 induces cardiac mesoderm differentiates into functional cardiomyocytes^[4].</p> <p>IWP-2 (GMP) (5 μM, treated at day 3) induces cardiomyocyte differentiation when applied following a pretreatment with Laduviglusib (GMP) (HY-10182G)^[5].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>RT-PCR^[3]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>hiPSCs</td> </tr> <tr> <td>Concentration:</td> <td>5 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>day 1 to 3</td> </tr> <tr> <td>Result:</td> <td>Reduced the expression of anti-cardiac mesoderm genes, and increased the expression of cardiac progenitors and cardiac genes (MYL2, TNNT3, and TNNT2).</td> </tr> </table>	Cell Line:	hiPSCs	Concentration:	5 μM	Incubation Time:	day 1 to 3	Result:	Reduced the expression of anti-cardiac mesoderm genes, and increased the expression of cardiac progenitors and cardiac genes (MYL2, TNNT3, and TNNT2).
Cell Line:	hiPSCs								
Concentration:	5 μM								
Incubation Time:	day 1 to 3								
Result:	Reduced the expression of anti-cardiac mesoderm genes, and increased the expression of cardiac progenitors and cardiac genes (MYL2, TNNT3, and TNNT2).								

CUSTOMER VALIDATION

- Adv Mater. 2021 Oct 10;e21104829.
- Dev Cell. 2020 Dec 21;55(6):679-694.e11.

- Clin Sci. 2023 Jan 13;137(1):109-127.
- Stem Cells Transl Med. 2021 May;10(5):743-755.
- Biochem Pharmacol. 2019 Nov;169:113608.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Guan J et al. Chemical reprogramming of human somatic cells to pluripotent stem cells. *Nature*. 2022;605(7909):325-331.
- [2]. Hamad S, et al. Generation of human induced pluripotent stem cell-derived cardiomyocytes in 2D monolayer and scalable 3D suspension bioreactor cultures with reduced batch-to-batch variations. *Theranostics*. 2019 Sep 25;9(24):7222-7238.
- [3]. Le MNT, et al. Auto/paracrine factors and early Wnt inhibition promote cardiomyocyte differentiation from human induced pluripotent stem cells at initial low cell density. *Sci Rep*. 2021 Nov 2;11(1):21426.
- [4]. Taniguchi J, et al. A synthetic DNA-binding inhibitor of SOX2 guides human induced pluripotent stem cells to differentiate into mesoderm. *Nucleic Acids Res*. 2017 Sep 19;45(16):9219-9228.
- [5]. Lian X, Zhang J, et al. Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/ β -catenin signaling under fully defined conditions. *Nat Protoc*. 2013 Jan;8(1):162-75.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA