



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Azido-PEG2-C2-amine

Cat. No.:	HY-140213
CAS No.:	166388-57-4
Molecular Formula:	C <sub>6</sub> H <sub>14</sub> N <sub>4</sub> O <sub>2</sub>
Molecular Weight:	174.2
Target:	PROTAC Linkers; ADC Linker
Pathway:	PROTAC; Antibody-drug Conjugate/ADC Related
Storage:	-20°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

#### In Vitro

Ethanol : 100 mg/mL (574.05 mM; Need ultrasonic)  
DMSO : 100 mg/mL (574.05 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	5.7405 mL	28.7026 mL	57.4053 mL
	5 mM	1.1481 mL	5.7405 mL	11.4811 mL
	10 mM	0.5741 mL	2.8703 mL	5.7405 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 90% corn oil  
Solubility: ≥ 2.5 mg/mL (14.35 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Azido-PEG2-C2-amine (N3-PEG2-CH2CH2NH2) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs<sup>[1]</sup>.

	Azido-PEG2-C2-amine is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs) [2]. Azido-PEG2-C2-amine is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.	
IC <sub>50</sub> & Target	PEGs	Non-cleavable Linker
In Vitro	<p>PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins<sup>[1]</sup>. ADCs are comprised of an antibody to which is attached an ADC cytotoxin through an ADC linker<sup>[2]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>	

## REFERENCES

- [1]. Su S, et al. Potent and Preferential Degradation of CDK6 via Proteolysis Targeting Chimera Degraders. J Med Chem. 2019 Aug 22;62(16):7575-7582.
- [2]. Tominari, Yusuk , et al. Heterocyclic compound. WO2020027225A1.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA