



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

linkedin.com/company/szaboscandic



Nrf2 activator-3

Cat. No.:	HY-143333		
CAS No.:	2766570-23-2		
Molecular Formula:	$C_{23}H_{18}F_3N_3O_2$		
Molecular Weight:	425.4		
Target:	Keap1-Nrf2		
Pathway:	NF-κB		
Storage:	Powder	-20°C 4°C	3 years 2 years
	In solvent	-80°C -20°C	6 months 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (235.07 mM; Need ultrasonic)

Preparing Stock Solutions	Concentration	Solvent Mass		
		1 mg	5 mg	10 mg
	1 mM	2.3507 mL	11.7536 mL	23.5073 mL
	5 mM	0.4701 mL	2.3507 mL	4.7015 mL
	10 mM	0.2351 mL	1.1754 mL	2.3507 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 90% corn oil
- Solubility: ≥ 2.5 mg/mL (5.88 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Nrf2 activator-3 is a potent Nrf2 activator. Nrf2 activator-3 is used for cerebral ischemic injury research ^[1] .	
In Vitro	<p>Nrf2 activator-3 (compound 24) ($1\text{ }\mu\text{M}$, $5\text{ }\mu\text{M}$, and $10\text{ }\mu\text{M}$) is against SNP ($400\text{ }\mu\text{M}$)-induced cell death with IC_{50} values of $76.86 \pm 3.54\text{ }\mu\text{M}$, $101.59 \pm 3.34\text{ }\mu\text{M}$, and $105.1 \pm 1.84\text{ }\mu\text{M}$ at $1\text{ }\mu\text{M}$, $5\text{ }\mu\text{M}$, and $10\text{ }\mu\text{M}$, respectively in PC12 cells^[1]¶</p> <p>Nrf2 activator-3 ($1\text{-}200\text{ }\mu\text{M}$) is against PC12 and hacat cell with IC_{50} values of $262.70 \pm 1.98\text{ }\mu\text{M}$ and $126.70 \pm 10.39\text{ }\mu\text{M}$, respectively^[1]¶</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Viability Assay^[1]</p>	
	Cell Line:	PC12 cell
	Concentration:	$1\text{ }\mu\text{M}$, $5\text{ }\mu\text{M}$, and $10\text{ }\mu\text{M}$

Incubation Time:	
Result:	Alleviated SNP-induced apoptosis in a concentration-dependent manner.

In Vivo

In the acute toxicity study, Nrf2 activator-3 (compound 24) shows toxicity to the experimental mice at 1000 mg/kg, the LD₅₀ of intraperitoneal injection is 789 mg/kg, and the 95% confidence interval was 550-1000 mg/kg in balb/c mice^[1].
 .In vivo pharmacokinetic properties study, Nrf2 activator-3 (5 mg/kg; Intraperitoneal injection) shows that plasma reached a maximum (323.06 ng/mL) at 2 h. the T_{max}, C_{max}, AUC_{0-inf}, F% and T_{1/2} values are 2 hour, 323.06 ng/mL, 2929.88 ng/mL*h, 28%, 12.75 hours respectively^[1].
 .Nrf2 activator-3 (5 mg/kg; i.v.) shows T_{max}, C_{max}, AUC_{0-inf}, and T_{1/2} values are 0.08 hours, 6911.14 ng/mL, 10182.73 ng/mL*h, and 8.26 hours respectively^[1].
 .Nrf2 activator-3 (3 mg/kg; 10 mg/kg;30 mg/kg) reduces the cerebral infarction volume and leads to decreased neurological deficits in MCAO rats^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	MCAO rats
Dosage:	3 mg/kg; 10 mg/kg; 30 mg/kg
Administration:	Intraperitoneal injection
Result:	Attenuated cerebral ischemic injury. (low dose: $16.37 \pm 6.51\%$, medium dose: $14.49 \pm 5.62\%$, high dose: $12.23 \pm 8.50\%$), which was similar to the effect of Edaravone ($12.77 \pm 5.82\%$).

Animal Model:	MCAO rats
Dosage:	3 mg/kg; 10 mg/kg; 30 mg/kg
Administration:	Intraperitoneal injection
Result:	Attenuated cerebral ischemic injury. (low dose: $16.37 \pm 6.51\%$, medium dose: $14.49 \pm 5.62\%$, high dose: $12.23 \pm 8.50\%$), which was similar to the effect of Edaravone ($12.77 \pm 5.82\%$).

REFERENCES

- [1]. Yaoqiang Lao, et al. Synthesis and biological evaluation of 1,2,4-triazole derivatives as potential Nrf2 activators for the treatment of cerebral ischemic injury. Eur J Med Chem. 2022 Jun 5;236:114315.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA