



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## MQA-P

Cat. No.:	HY-149203
Molecular Formula:	C <sub>40</sub> H <sub>36</sub> BrN <sub>2</sub> O <sub>2</sub> P
Molecular Weight:	687.6
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



## BIOLOGICAL ACTIVITY

<b>Description</b>	<p>MQA-P is a multifunctional near-infrared (NIR) fluorescent probe for simultaneously detecting ONOO<sup>-</sup>, viscosity, and polarity within mitochondria. MQA-P exhibits a remarkable turn-on response to ONOO<sup>-</sup> (<math>\lambda_{em}</math>=645 nm) and is highly sensitive to viscosity/polarity in the NIR channel with <math>\lambda_{em}</math>&gt;704 nm. MQA-P exhibits excited-state intramolecular charge transfer (ESICT) feature that is highly polarity-sensitive by engineering N,N-dimethylamino as the electron donor and a quinoline cationic unit as the electron acceptor. MQA-P is used for ferroptosis or cancer diagnosis in vitro and in vivo via dual-channel images [1].</p>
<b>In Vitro</b>	<p>Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs).</p> <ol style="list-style-type: none"> <li>MQA-P is dissolved in dimethyl sulfoxide (DMSO) to prepare a stock solution (1.0 mM).</li> <li>For imaging of ONOO<sup>-</sup> in live cells. HeLa cells are incubated with MQA-P (5 <math>\mu</math>M) for 30 min as control; pretreated with SIN-1 (HY-126849; 100 <math>\mu</math>M) for 30 min and then incubated with MQA-P (5 <math>\mu</math>M) for another 30 min. The fluorescence images are obtained on a confocal laser scanning microscope with a green channel (<math>\lambda_{ex}</math>= 405nm, <math>\lambda_{em}</math>= 550-670 nm).</li> <li>For imaging of viscosity in live cells. HeLa cells were incubated with MQA-P (5 <math>\mu</math>M) for 30 min as control; pretreated with Monensin (HY-N4302; 10 <math>\mu</math>M) for 30 min and then incubated with MQA-P (5 <math>\mu</math>M) for another 30min. The fluorescence images are obtained on a confocal laser scanning microscope with a red channel (<math>\lambda_{ex}</math>= 561 nm, <math>\lambda_{em}</math>= 680-750 nm).</li> <li>For dual-channel imaging of ONOO<sup>-</sup>, viscosity and polarity during ferroptosis. HeLa cells are incubated with MQA-P (5 <math>\mu</math>M) for 30 min as control; pretreated with Erastin (HY-15763; 50 <math>\mu</math>M) for 30 min and then incubated with MQA-P (5 <math>\mu</math>M) for another 30 min. The fluorescence images are obtained on a confocal laser scanning microscope with a green channel (<math>\lambda_{ex}</math>= 405nm, <math>\lambda_{em}</math>= 550-670 nm) for ONOO<sup>-</sup> and a red channel (<math>\lambda_{ex}</math>= 561 nm, <math>\lambda_{em}</math>= 680-750 nm) for viscosity and polarity<sup>[1]</sup>.</li> </ol> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
<b>In Vivo</b>	<p>Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs).</p> <ol style="list-style-type: none"> <li>For tissue slices imaging, the normal organs (including heart, liver, spleen, lung, and kidney) and tumor are isolated from the mice, then sectioned as 5 <math>\mu</math>m thicknesses, respectively.</li> <li>These slices are incubated with MQA-P (20 <math>\mu</math>M) for 30 min, then washed with PBS (pH 7.4) three times, and finally subjected to in vivo imaging using a confocal laser scanning microscope with a green channel (<math>\lambda_{ex}</math>=405nm, <math>\lambda_{em}</math>=550-670 nm) for ONOO<sup>-</sup> and a red channel(<math>\lambda_{ex}</math>=561 nm, <math>\lambda_{em}</math>=680-750 nm) for viscosity and polarity, respectively<sup>[1]</sup>.</li> </ol> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

---

## REFERENCES

[1]. Li Fan, et al. Multifunctional Fluorescent Probe for Simultaneous Detection of ONOO-, Viscosity, and Polarity and Its Application in Ferroptosis and Cancer Models. Anal Chem. 2023 Apr 4;95(13):5780-5787.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA