



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Imepitoin

<b>Cat. No.:</b>	HY-14953		
<b>CAS No.:</b>	188116-07-6		
<b>Molecular Formula:</b>	C <sub>13</sub> H <sub>14</sub> ClN <sub>3</sub> O <sub>2</sub>		
<b>Molecular Weight:</b>	279.72		
<b>Target:</b>	GABA Receptor		
<b>Pathway:</b>	Membrane Transporter/Ion Channel; Neuronal Signaling		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 12.5 mg/mL (44.69 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	3.5750 mL	17.8750 mL	35.7500 mL
		5 mM	0.7150 mL	3.5750 mL	7.1500 mL
10 mM		0.3575 mL	1.7875 mL	3.5750 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	<ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: ≥ 1.25 mg/mL (4.47 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.25 mg/mL (4.47 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: ≥ 1.25 mg/mL (4.47 mM); Clear solution</li> </ol>				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Imepitoin (AWD 131-138) is a new low-affinity partial benzodiazepine receptor agonist with potent anticonvulsant and anxiolytic properties in rodent models.
<b>IC<sub>50</sub> &amp; Target</b>	GABA receptor
<b>In Vitro</b>	AWD 131-138 dose-dependently stimulated GABA currents(Recombinant gamma-aminobutyric acid A (GABA(A)) receptors of the subunit compositions alpha1beta2gamma2, alpha1beta3gamma2, alpha2beta2gamma2, alpha3beta2gamma2 and

---

alpha5beta2gamma2). At 10 microM AWD 131-138, this allosteric stimulation amounted in average to about 12-21% of the maximal stimulation achieved using diazepam. The threshold of stimulation was about 0.3-1.0 microM [1].  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

**In Vivo**

AWD 131-138 did not produce midazolam-like responding or alter response rates at cumulative doses up to 18.0 mg/kg i.m. (plasma levels over 2100 ng/ml). When AWD 131-138 (10-100 microg/kg/injection) was studied by substitution, responding declined to vehicle substitution levels within three sessions. At the dose of 100 microg/kg i.v. AWD 131-138, sufficient drug was self-administered during the first session (about 3.5 mg/kg) to produce plasma levels above 1000 ng/ml, yet responding over the next two sessions dropped to vehicle levels [2]. Prolonged oral administration with twice-daily dosing of ELB 138 with either 5 or 40 mg/kg over a 5-week period was not associated with loss of anticonvulsant efficacy in the PTZ dog model [3].  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## REFERENCES

- [1]. Sigel E, et al. The antiepileptic drug AWD 131-138 stimulates different recombinant isoforms of the rat GABA(A) receptor through the benzodiazepine binding site. *Neurosci Lett.* 1998 Apr 3;245(2):85-8.
- [2]. Yasar S, et al. Evaluation of the novel antiepileptic drug, AWD 131-138, for benzodiazepine-like discriminative stimulus and reinforcing effects in squirrel monkeys. *Eur J Pharmacol.* 2003 Apr 4;465(3):257-65.
- [3]. Loscher W, et al. Anticonvulsant efficacy of the low-affinity partial benzodiazepine receptor agonist ELB 138 in a dog seizure model and in epileptic dogs with spontaneously recurrent seizures. *Epilepsia.* 2004 Oct;45(10):1228-39.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA